

# Influence of microbial probiotics on ruminant health and nutrition: sources, mode of action and implications

Rine C Reuben,<sup>a</sup>  Mona M M Y Elghandour,<sup>b</sup>  Othman Alqaisi,<sup>c</sup>  
John W Cone,<sup>d</sup> Ofelia Márquez<sup>e</sup> and Abdelfattah Z M Salem<sup>b\*</sup> 

## Abstract

Globally, ruminant production contributes immensely to the supply of the highest quality and quantity of proteins for human consumption, sustenance of livelihoods, and attainment of food security. Nevertheless, the phasing out of antibiotics in animal production has posed a myriad of challenges, including poor growth, performance and nutrient utilization, pathogen colonization, dysbiosis, and food safety issues in ruminants. Probiotics (direct-fed microbials), comprising live microbial strains that confer health and nutritional benefits to the host when administered in appropriate quantities, are emerging as a viable, safe, natural and sustainable alternative to antibiotics. Although the mechanisms of action exerted by probiotics on ruminants are not well elucidated, dietary probiotic dosage to ruminants enhances development and maturation, growth and performance, milk production and composition, nutrient digestibility, feed efficiency, pathogen reduction, and mitigation of gastrointestinal diseases. However, the beneficial response to probiotic supplementation in ruminants is not consistent, being dependent on the microbial strain selected, combination of strains, dose, time and frequency of supplementation, diet, animal breed, physiological stage, husbandry practice, and farm management. Nonetheless, several studies have recently reported beneficial effects of probiotics on ruminant performance, health and production. This review conclusively re-iterates the need for probiotics inclusion for the sustainability of ruminant production. Considering the role that ruminants play in food production and employment, global acceptance of sustainable ruminant production through supplementation with probiotics will undoubtedly ensure food security and food safety for the world.

© 2021 Society of Chemical Industry.

**Keywords:** probiotics; nutrition; ruminant; health; food security

## INTRODUCTION

Apart from the production of high-quality protein in sufficient quantities for human consumption, sustenance of livelihoods (especially amongst rural dwellers) and contribution to food security comprise the other important roles of animal husbandry and production.<sup>1</sup> Livestock also provide draught animal power to millions of disadvantaged, marginal and smallholder farmers, mostly in developing countries, for both cultivation of farmlands and small-scale transport. Also, the manure produced by these animals is often used as biofuel and biofertilizer.<sup>2,3</sup> With the projected increase in the global population to nine billion people in 2050<sup>4</sup> and global economic growth and urbanization,<sup>5</sup> the increasing demand for livestock products, including meat, milk and eggs, is putting immense pressure on the livestock industry to continuously meet global expectations in production despite scarce resources.<sup>2</sup> There is a need for the development of a sustainable global animal production that will make the most efficient use of limited resources for the production of food and conserving the environment.

The importance of ruminants in the production of high-quality protein (meat and milk) and their ability to efficiently utilize fibrous feeds cannot be overemphasized. In comparison with other livestock species, ruminants are the primary food source

for humans, especially in the developing parts of the world most affected by desertification and scarce fertile agricultural soils.<sup>6</sup> Furthermore, ruminants occupy the largest land area globally and have considerable influence on the environment through their emission of greenhouse gases, including carbon dioxide, methane and nitrous oxide.<sup>7,8</sup> Over the years, antibiotics have

\* Correspondence to: A Z M Salem, Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma del Estado de México, Toluca, 50000, Mexico. E-mail: asalem70@yahoo.com; salem@uaemex.mx

a German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Leipzig University, Leipzig, Germany

b Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma del Estado de México, Toluca, Mexico

c Animal and Veterinary Sciences Department, College of Agricultural and Marine Sciences, Sultan Qaboos University, Muscat, Oman

d Animal Nutrition Group, Wageningen University & Research, Wageningen, The Netherlands

e Centro Universitario UAEM Amecameca, Universidad Autónoma del Estado de México, Amecameca, Mexico

been used in the livestock industry at both prophylactic and sub-therapeutic levels to enhance animal production (growth performance) and also to mitigate infectious diseases.<sup>9</sup> Nevertheless, with the phasing out of antibiotics in animal production, as a result of immense public health concerns, including the presence of drug residues in animal products,<sup>1,10,11</sup> the emergence and spread of resistance, and dysbiosis of gut microflora, there is a need for the application of naturally safe alternatives that will both improve animal growth performance as well as mitigate infectious diseases.

Probiotics or direct-fed microbials (DFM) are now widely accepted as safe and sustainable alternatives to antibiotics in animal production. The significant effects of probiotic supplementation as alternatives to antibiotics in animal production have been reported, as reviewed previously.<sup>12</sup> Probiotics can act by ameliorating enteric infections through the competitive exclusion of pathogens, mitigating chronic inflammatory and allergic diseases via immunomodulation and immune-stimulation, increasing digestibility and nutrient assimilation, improving the intake of dry matter to feed conversion ratio, reducing the emission of greenhouse gases, promoting growth and health performance, and improving ruminant meat and milk production.<sup>12-16</sup> The activities of probiotics can vary and mostly depend on both intrinsic and extrinsic factors, including the specific probiotic strain in use, dose, host species, probiotic strain combination (for multi-strain probiotics), environmental factors, husbandry practices and farm management.<sup>1</sup>

This review examines current knowledge and the concept of probiotic supplementation in ruminant production. It further explores the mechanisms of action, as well as the specific influence that probiotics exert on ruminants, including development and maturation, growth and performance, milk production and composition, nutrient digestibility and feed efficiency, haemato-biochemical parameters, pathogen reduction, and mitigation of gastrointestinal diseases. The review also discusses the sources, features and administration of ruminant probiotics, and, finally, the inclusion of probiotics in the sustainability of ruminant production as the biological strategy for global food security and food safety.

## PROBIOTICS: CONCEPT AND STRAINS

### Concept of probiotics and their strains

Over the years, researchers have proposed many definitions for the term "probiotic" based on the beneficial and health outcomes obtained as a result of the application of certain microbial strain(s) in different host species. Kollath, in 1953,<sup>17</sup> initially used the term "probiotic" to generally describe inorganic and organic substances that were considered to restore the health of malnourished patients. Lilly and Stillwell, in 1965,<sup>18</sup> further expanded the term "probiotic" to mean "unknown substances" produced by certain ciliate protozoa, which enhanced the survival and development of other protozoa. Subsequently, Parker, in 1974,<sup>19</sup> further included both living organisms and non-living substances in his definition of probiotics to be "organisms and substances which contribute to intestinal microbial balance". The increasing application of probiotics in feed supplementation in livestock production led to the reconsideration of probiotics as "alive microbial feed supplement which beneficially affects the host animal by improving its intestinal microbial balance".<sup>20</sup> Other researchers defined probiotics as "live cultures of microorganisms that are deliberately introduced into the rumen to improve animal health

or nutrition".<sup>21</sup> With the massive production and commercialization of probiotics for livestock production, the Food and Drug Administration of the United States requires manufacturers of livestock feeds to replace the term "probiotic" with the phrase "direct-fed microbial" (i.e. DFM).<sup>22</sup> DFM has been defined as "alive, naturally occurring microorganisms that have been used to improve the digestive function of livestock".<sup>23,24</sup>

The broader perspectives in the definition and concept of DFM included a broad spectrum of microorganisms, such as bacteria, yeast, fungi, fragments of microbial cells and microbial secretions.<sup>25-29</sup> Microbial strains used as DFM flourish in the rumen of livestock and modulate the microflora and fermentation activities to benefit the host.<sup>30</sup> However, the increasing interest generated in probiotic science and a progressive understanding of probiotic action led to modification of the definition of a probiotic. The current and widely accepted definition of the term probiotic was proposed by a joint Food and Agriculture Organization and World Health Organization working group, who defined probiotics as "live microbial strains that, when administered in adequate amounts, confer a health benefit on the host".<sup>31</sup>

In ruminant production, the specific probiotics used as DFM include different bacterial and fungal species such as *Bacillus*, *Bifidobacterium*, *Enterococcus*, *Lactobacillus*, *Propionibacterium*, *Megasphaera elsdenii* and *Prevotella bryantii*, as well as strains of *Aspergillus* and *Saccharomyces*, respectively.<sup>30,32</sup> Nevertheless, strains of lactic acid bacteria (LAB) including *Lactobacillus*, *Bifidobacterium* and *Streptococcus*, have gained predominance and are mostly the widely used species as probiotic strains in the supplementation of ruminant feed.<sup>33</sup> It is noteworthy to add that many of the microbial strains used as ruminant probiotics exist naturally in the gastrointestinal tract of healthy animals but are often disrupted by both intrinsic and extrinsic stress factors on the host species.<sup>33</sup>

Ruminant probiotic products are usually comprised of mono- and/or multi-strains or multi-species. Also, allochthonous microorganisms, such as yeasts which are normally absent in the ruminant gastrointestinal tract (GIT) are used as probiotics, mostly in combination with microbial species such as *Bifidobacterium* and *Lactobacillus*, which are regarded as indigenous inhabitants of the ruminant gut.<sup>2</sup> Probiotics mainly composed of bacteria are effective in young ruminants, whereas studies have shown that fungal probiotics exert their beneficial effects mostly in adult ruminants.<sup>34</sup>

### Ideal probiotic characteristics

There are many criteria for the selection of probiotics. First, microbial strains to be used as probiotics must successfully pass different *in vitro* evaluations before they can be approved for *in vivo* use. An effective and ideal probiotic is expected to function efficiently under a wide variety of conditions, including industrial processes and also in the gastrointestinal tract. There are a variety of parameters used for the screening and evaluation of probiotics, depending on the intended purpose and application of the microbial strain(s) in the target host(s).<sup>35,36</sup>

The initial and major step in the choice of a potential probiotic strain is the identification of its taxonomic class, which may indicate the domain and physiology of the specific strain. These features have immense effects on the choice of potential and novel probiotic strains.<sup>37</sup> Nevertheless, the effectiveness exhibited by a microbial strain intended for use as a probiotic is more important than the origin or source of such a strain.<sup>31</sup> Also, because a single probiotic strain is unlikely to possess all the equally suitable health

benefits, combining suitable multiple strains with synergistic effects is often better.<sup>38</sup>

The microbial strain(s) to be used as probiotic either for humans or animals must be (i) identified to species and strain levels; (ii) examined for antimicrobial susceptibility; (iii) characterized for antimicrobial production; (iv) examined for toxicity and pathogenicity; and (v) compatible with other authorized additives (where necessary).<sup>39</sup>

### Probiotic sources and administration in ruminants

The source and origin of probiotic strains are directly dependent on the intended use and application of such probiotics. Generally, probiotics can be of animal or human origin or from food sources such as raw milk or fermented foods.<sup>40</sup> It was reported that the microflora of neonatal animals reflects the bacterial composition that they consumed primarily from fresh maternal milk. As such, ruminants' milk may be a repository of natural microbiota and could be used primarily for probiotics isolation. Furthermore, microbial strains intended to be used as probiotics should be appropriately isolated and identified before use.<sup>41,42</sup> Nevertheless, a report on probiotics from the Food and Agriculture Organization<sup>31</sup> rated the specificity and functionality of probiotic action above the source of probiotic microorganisms.

Probiotics can be administered through different routes. However, oral and vagina routes are often used.<sup>43</sup> The major route for the administration of probiotics in ruminants is via the oral cavity.<sup>44-48</sup> Additionally, recent studies report probiotics administration through the vaginal of ruminants. For example, Deng *et al.*<sup>49</sup> administered a mixture of lactic acid bacteria (LAB) probiotics to periparturient cows through intravaginal infusion.

## MODE OF PROBIOTIC ACTION IN RUMINANTS

The advent of high-throughput molecular techniques has made it easier to obtain a better understanding of the diversity and mechanisms of action of probiotic strains in their host. The detailed mechanisms of probiotic action in ruminants have not been clearly and fully understood to date. However, several studies have illustrated some major processes used by probiotics in exerting their benefits on ruminants.<sup>44,45,50,51</sup> These mechanisms are specific to bacterial and fungal probiotics or their combinations.

### Bacterial probiotics

As previously stated, strains of *Bifidobacterium* and *Lactobacillus* spp. are the major bacterial probiotics used in ruminant diets. The mode of action of these bacterial strains used as ruminant probiotics solely depends on the specific strain(s) of bacteria selected, dosages, frequencies and times of supplementation, and, to some extent, farm management. Although some probiotic strains act mainly with the rumen, other specifically impact the GIT of ruminants.<sup>52</sup>

#### Bacterial probiotic action within the ruminant rumen

Although not well elucidated, the action of probiotics within the rumen primarily depends on the LAB and lactic acid utilizing bacteria (LUB).<sup>53</sup> LAB produce organic acids that beneficially mitigate acidosis within the rumen, most especially in dairy cows by stimulating the growth of beneficial microbes and LUB.<sup>29,54</sup> LUB act by decreasing lactic acid concentration within the rumen hence, maintaining a steady pH in the rumen. When ruminants feed on fermentable diets, lactate is produced in large amounts in their

rumen.<sup>55</sup> Certain bacteria, such as *Megasphaera elsdenii*, utilize the accumulated lactate, thereby impeding acidosis, which may be detrimental to the animal.<sup>56</sup> Ruminants harbor *Propionibacteria* in great numbers within their rumen, and these bacteria efficiently act on ruminant diets that are composed mainly of forage and concentrate.<sup>55</sup> Strains of *Propionibacteria* improve fermentation with a consequential increase in the production of propionate in the rumen.<sup>57</sup> During ruminal fermentation in early lactation, *Propionibacteria* convert lactate to propionate,<sup>55</sup> which leads to an increased production of liver glucose, making substrates available for lactose synthesis, hence reducing ketosis at the same time as improving energy efficiency.<sup>57,58</sup> Propionate accounts for the release of approximately 61–67% of glucose in lactating cows and growing ruminants.<sup>59</sup> The decrease in the emission of greenhouse gas, especially methane, by ruminants has been reported to be the result of an increase in ruminal propionate.<sup>60</sup> Although not well understood, Ghorbani *et al.*<sup>61</sup> reported that the inclusion of *Propionibacteria* strains as DFM in ruminants' diets decreased the numbers of amylolytic bacteria at the same time as increasing the population of *Entodinium* (protozoa) in the rumen of feedlot steers.

The overall efficiency and functionality of bacterial strains used as ruminant probiotics, either as mono- and/or multi-strains or species, can mostly be determined by their effect on the microflora in the rumen.<sup>62,63</sup> Furthermore, major environmental factors, including the quantity of lactic acid and volatile fatty acids (VFA), as well as pH, also determine the survival of microbes in the rumen.<sup>64,65</sup>

#### Bacterial probiotic action within post-ruminal GIT

Over the years, researchers have extensively explored the benefits exerted by bacterial probiotics and their mechanisms of action on the post-ruminal GIT.<sup>30</sup> The major modes of probiotic action within the post-ruminal GIT are as discussed below.

- (1) Competitive exclusion of pathogens from adhering to intestinal mucosa. One of the mechanisms considered to be the most important and beneficial, which is exhibited by probiotic bacteria, is the competitive exclusion of pathogens.<sup>66</sup> This is based on the ability of bacteria to interact with each other and compete for nutrients and adhesion sites on the intestinal mucosa. Beneficial bacteria gain competitive advantage by modifying and rendering their microenvironment unsuitable for their competitors, mostly pathogens.<sup>67</sup> Frizzo *et al.*<sup>68</sup> showed how LAB used as probiotics adhered to the gut and protected animals against *Salmonella*. Also, the adhesion of enteropathogenic *Escherichia coli* and enterotoxigenic *Klebsiella pneumoniae* to intestinal mucosa were significantly reduced through the inclusion of *Lactobacillus rhamnosus* (Lcr35) as DFM.<sup>69</sup>
- (2) Secretion of antimicrobial substances. Once probiotic strains are established within the gut, they can synthesize substances with either bacteriostatic or bactericidal properties such as bacteriocins, hydrogen peroxide and short chain fatty acids, as well as several organic acids. These substances secreted by probiotics, as well as the lowered intestinal pH, are detrimental and harmful to intestinal pathogens.<sup>70</sup> Bacteriocins antagonize pathogens within the GIT by interfering with DNA replication of target pathogens.<sup>71</sup> Most bacterial species, including *Bacillus* spp., LAB and bifidobacteria used as probiotics,<sup>72,73</sup> are capable of producing several types of thermostable bacteriocins<sup>74</sup> with a wide range of antimicrobial

activity against potential animal pathogens including *Bacillus cereus*, *Staphylococcus*, *Enterococcus*, *Listeria* and different species of *Salmonella*.<sup>75</sup>

- (3) Immunomodulation and immune-stimulation. The immune system component of the gut, which protects the host from different types of antigens in the lumen of the GIT, can be affected by probiotics. Probiotic action within the intestinal tract modulates both the innate and adaptive immunity of the animal.<sup>2</sup> The ability of probiotics to adhere to the intestinal mucosa creates a natural barrier against potential pathogens and, in some instances, enhances the host immunity. Probiotics stimulate the immune system by increasing the synthesis and secretion of immunoglobulins, as well as macrophage and lymphocyte activities, and also by stimulating  $\gamma$ -interferon production.<sup>24</sup> *Lactobacillus casei* Shirota and *L. rhamnosus* Lr23 are known probiotics strains that stimulate macrophages to secrete tumor necrosis factor- $\alpha$  when used in ruminants.<sup>76</sup> Although the pro-inflammatory effect exerted by probiotics could positively impact the animal by mitigating potential pathogens from the GIT, it could also pose a negative impact by creating a pro-inflammatory environment that may damage the entire animal.
- (4) Colonization resistance. Microorganisms colonizing the GIT of neonatal ruminants generally originate from the adult. These young ruminants become protected against intestinal pathogens by these colonizing microorganisms. Nevertheless, the intensive ruminant production has reduced this natural microbial colonization of the gut, hence rendering animals more susceptible to pathogen challenges in the GIT. In neonates, probiotics could mimic natural colonization, or colonize adult animals, thus shielding the intestinal mucosa from colonization by pathogenic organisms.<sup>2</sup> The presence of hydrophobic surface layer proteins in most probiotic strains, especially *Bifidobacterium* and *Lactobacillus* spp., enables them to adhere to the surface of their host enterocytes.<sup>77</sup> These adhering probiotic strains bind to certain receptor sites on the intestinal epithelium thereby preventing pathogens from accessing and colonizing intestinal epithelial cell surfaces.<sup>77,78</sup>
- (5) Alteration in pathogenicity and virulence gene expression of pathogens. Cell to cell communication in bacteria through auto-inducers (secretory chemical signals) affects bacterial physiology.<sup>79</sup> Quorum sensing, one process of bacterial communication, is also used for communication between bacteria and their host.<sup>80</sup> Through quorum sensing, probiotics can influence the virulence and pathogenicity of pathogens. The ability of enterohaemorrhagic *E. coli* serotype O157:H7 to secrete extracellular autoinducer-2 was reported to be substantially inhibited by fermentation products from *Lactobacillus acidophilus* La-5, which suppressed the expression of the locus of enterocyte effacement (LEE) virulence gene in *E. coli* serotype O157:H7. Consequently, quorum sensing can become disrupted so that colonization and pathogenesis by *E. coli* serotype O157:H7 in the GIT is prevented.<sup>81</sup> Using experimental rats, *Lactobacillus salivarius* decreased the expression of *Listeria monocytogenes* virulence gene, listeriolysin O in intestinal villi and Peyer's patches. Also, both *Lactococcus lactis* and *L. salivarius* lowered the *Listeria* count in spleens.<sup>82</sup> Likewise, Dong et al.<sup>83</sup> reported repression in the expression of three virulence genes of *L. monocytogenes*, including InIA, InIB and prfA, after treatment with *Lactobacillus plantarum* CICC6257.

### Fungal and yeast probiotics

Because of the role of fungi in the stimulation of ruminal fermentation, fungal probiotics have been used extensively in ruminants.<sup>53</sup> Strains of *Saccharomyces* and *Aspergillus* are the most common fungal probiotics used in ruminants.<sup>52</sup> The inclusion of fungal-based probiotics in ruminant diets stabilizes the pH of the rumen, with a significant increase in total VFAs and a reduction in the concentration of ammonia.<sup>26,84,85</sup> The presence of yeast in the rumen also stimulates the enzymatic activity and growth of cellulolytic bacteria, in addition to improving the digestion of high-fiber diets and microbial translation of mRNA to proteins.<sup>86</sup> The DFM have been shown to reduce the redox potential in the rumen, thereby creating a desirable environment for the growth and survival of strict anaerobic organisms that often secrete vitamins, fatty acids and other factors stimulating ruminal microbial biomass synthesis.<sup>87,88</sup> The use of yeast DFM has been shown to mitigate rumen acidosis, as well as enhance the activity of LUB, especially *Selenomonas ruminantium*.<sup>89</sup> When used as ruminant probiotics, yeast mainly acts by improving the digestion of high fiber and concentrate diets, enhancing feed efficiency and also stimulating rumen fermentation.<sup>53,84</sup> More work is needed to fully unravel the detailed mechanisms of action used by fungal-based probiotics in improving ruminant production.

## PROBIOTICS IN RUMINANT NUTRITION

### Effects of probiotics on ruminants

The application of probiotics in ruminants has yielded significant results, especially with respect to boosting their health status, as well as enhancing productivity and their general performance. Researchers have concentrated more on the application of ruminant probiotics in the pre-ruminant's life and, to some extent, in adult ruminants. Over the years, studies have centered mostly on aspects of ruminants' health, including the control of pathogen carriage, infectious disease mitigation (mostly enterodiseases and mastitis), reduction of the severity of diarrhoea, and economic parameters such as the reduction of feed intake and feed conversion ratio. The dominant ruminant species investigated for probiotic applications are calves, goats and, to some extent, lambs/ewes.

#### Development and maturation of ruminants

With the application of dietary supplementation of probiotics in ruminants, several microbial strains, including *Lactobacillus*, *Bifidobacterium*, *Enterococcus*, *Streptococcus* and *Propionibacterium* spp., have been successfully used for promoting the health, development and maturation of juvenile ruminants.<sup>23,68,71,90-92</sup> The enhancement of growth, development and maturation of neonatal ruminants as a result of probiotic supplementation has gained increased merit in animal production.<sup>91</sup> Neonatal calves are highly vulnerable to pathogens before and during weaning. During weaning, calves become highly susceptible to pathogen colonization in their gut as a result of their immature gut microbiota and a sudden change in feed composition.<sup>25,93</sup> The rapid adaptation to the change in feed composition and rapid colonization of beneficial microorganisms within the GIT of calves is of immense importance in ruminant maturation and development.<sup>23</sup> This crucial developmental stage prevents the colonization of enteropathogens within the GIT of calves. These enteropathogens are responsible for diarrhoea in calves that can result in a huge economic costs.<sup>23</sup> Apart from the impact of colostrum (maternal first milk) in ruminants, the introduction of probiotics into the immature

GIT of young ruminants can promote health.<sup>91</sup> At birth, microbial colonization of the gut increases, and the maturation of gut microflora and the intestinal mucosa proceeds gradually.<sup>94</sup> Shortly after birth, calves receive beneficial microorganisms, such as *Lactobacillus* spp., *Enterococcus* spp. and non-pathogenic *E. coli* before the incorporation of other microbial strains (mainly anaerobes) into the forestomach gut.<sup>92</sup> Maternal milk, maternal gut microflora and the environment are the primary sources of probiotic microorganisms for neonatal calves.<sup>91,94</sup> Maternal milk contains antibodies, lactoferrin,  $\kappa$ -casein, oligosaccharides and other proteins that stimulate the growth of *Bifidobacterium* spp. and other probiotic microorganisms within the gut and also antagonize potential pathogens.<sup>95</sup> The growth of these beneficial microorganisms in the gut facilitates the ability of young ruminants to digest high fiber diets at the same time as supporting their overall development and maturation<sup>91</sup> (Table 1).

#### Growth and performance

The supplementation of ruminants with specific mono- and/or multi-strain probiotics has been shown to significantly increase their growth and overall performance, including daily body weight, body weight gain and feed efficiency.<sup>23,30,96</sup> Although the effects of probiotics on the growth and performance of ruminants may vary, scientific investigations suggest that an increase and modulation of the intestinal microflora, as well as stimulation of the activities of cellulolytic bacteria that enhance digestion and absorption of nutrients, comprise the main modes of action exhibited by probiotics aimed at promoting growth and performance.<sup>25,85,97-99</sup>

The supplementation of cattle, goats and sheep with probiotics has led to better performance, improved feed intake and efficiency and weight gain.<sup>44,45,50,57,96,100</sup> The inclusion of *Pediococcus acidilactici* and *Pediococcus pentosaceus* in the diet of lambs during the post-weaning period significantly improved their average daily weight gain (+25.2 g per lamb), final body weight (+3.16 kg) and feed conversion ratio (FCR, -1.18).<sup>101</sup> However, the supplementation of small ruminants with probiotics can improve FCR.<sup>56</sup> Dietary inclusion of live yeast in beef cattle and dairy cows reportedly improved FCR, average daily gain, feed intake and final weight.<sup>25,57,100,102</sup> The inclusion of multi-species probiotics, containing *Lactobacillus*, *Streptococcus*, *Aspergillus* and *Saccharomyces cerevisiae*, in the diet of growing lambs significantly increased the average daily weight gain by 7.2%.<sup>103</sup> Moreover, Mudgal and Baghel<sup>104</sup> reported a 31.4% increase in the average daily weight gain of buffalo calves fed with diets containing *L. acidophilus*.

Nevertheless, some studies recently reported divergent findings, showing that no effect was exerted on the growth and performance of ruminants supplemented with probiotics.<sup>45,46,50</sup> The difference in the microbial strains used as probiotics, the number of cells used (dose), ruminant species and breed, environmental factors, feeding practice, farm management, and other variables may account for their contrary results.<sup>30,96</sup> As shown in Table 1, probiotic effects are species-specific and host-dependent.<sup>43,105,106</sup>

#### Milk composition and milk production

The inclusion of probiotics in the feeding of livestock may not only improve health status, but may also ensure food security and food safety.<sup>107</sup> The supplementation of ruminants with probiotics has been shown to increase milk production, milk quality and milk composition, with a higher presence of functional components

such as protein, fat and solid non-fat and a lower count of somatic cells.<sup>23,33,100,108,109</sup> The inclusion of *S. cerevisiae* and *Aspergillus oryzae* in the diets of dairy cows significantly increased milk production, with a greater concentration of milk proteins.<sup>110</sup> Ruminal fermentation and milk production also improved when dairy cows were supplemented with strains of *Bacillus subtilis* as probiotics.<sup>111,112</sup> Also, Stella *et al.*<sup>113</sup> reported a 14.0% increase in the average daily milk yield among lactating Saanen dairy goats fed with  $4 \times 10^9$  cfu day<sup>-1</sup> per animal of *S. cerevisiae*.

The overall findings of a meta-analysis on the effects of a *S. cerevisiae* probiotic strain on milk production, feed intake and the rumen fermentation of buffaloes, cattle, goats and sheep showed a significant increase in milk yield.<sup>114</sup> Similarly, there was an approximate 8.0% increase (i.e. 1.1 kg day<sup>-1</sup>) of milk yield by Tunisian Holstein Friesian cows supplemented with yeast probiotics compared to the control.<sup>115</sup> From the same experiment, the probiotic supplemented cows had higher protein (41.7 g per cow per day) and milk fat (53 g per cow per day) levels than the control animals (38.7 and 47 g per cow per day). Different mechanisms have been reported to explain the increases in milk yield in response to probiotic supplementation. On the one hand, it has been reported that increases in milk yield result from the ability of probiotic strains to reduce inflammation of the udder and thus suppress the number of somatic cells in milk.<sup>109</sup> On the other hand, other studies have postulated that improvements in milk production and composition are the result of probiotics with respect to stimulating growth and an increase in the population of both fiber-degrading and cellulolytic-degrading bacteria<sup>116</sup> (Table 1).

#### Feed efficiency and nutrient digestibility

Improvement in ruminant growth and overall performance as a result of probiotic supplementation is often associated with improvement in fiber degradation, feed intake, nutrient digestibility and ruminal digestion.<sup>23,117-119</sup> As a complex carbohydrate and usually non-digestible by mammalian enzymes, fiber, being the major component of plants, is mainly composed of cellulose, hemicellulose and lignin.<sup>120,121</sup> Between 15% and 70% of diets fed to ruminants are composed of cellulose and hemicellulose.<sup>122</sup> The ability of probiotics to improve nutrient digestibility and feed intake in ruminants may be attributed to the enhanced proliferation and growth of ruminal cellulolytic bacteria and the prevention of acidosis in the rumen.<sup>123</sup> The supplementation of lactating cows with yeast increased the frequency of feed intake, fiber degradation and nutrient digestibility in the rumen.<sup>25</sup> Feeding Awassi lambs with yeast culture has been reported to increase the digestibility of dry matter (676 g kg<sup>-1</sup>), crude protein (653 g kg<sup>-1</sup>), organic matter (683 g kg<sup>-1</sup>) and neutral detergent fiber (which is the most common measure of fiber used for animal feed analysis) (561 g kg<sup>-1</sup>) compared to the control.<sup>124</sup> Digestion of acid and neutral detergent fiber, as well as levels of crude protein, improved after lactating Holstein cows were supplemented with a diet containing strains of *Lactobacillus acidophilus* NP51 and *Propionibacterium freudenreichii* NP24.<sup>125</sup> Azzaz *et al.*<sup>126</sup> reported improved digestibility of organic matter, dry matter, crude fiber, crude protein and nitrogen-free extract in goats fed with diets containing *L. acidophilus* and *Aspergillus awamori*. The inclusion of dried live yeast significantly improved the digestibility of fiber, dry matter and crude protein in lambs fed diets comprised of concentrates and roughages at a 60:40 ratio in the dry matter.<sup>127</sup> Nevertheless, variations in probiotic species and strains, dosage, feed composition, feeding systems and

**Table 1.** Summary of probiotics effects on ruminant nutrition and performance

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
Bio-Plus 2B®, Chr. (Hansen A/S)	<i>Bacillus subtilis</i> , <i>B. licheniformis</i>	132 × 10 <sup>9</sup> cfu kg <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased growth rate</li> <li>2. Increased feed intake</li> <li>3. Increased average daily gain</li> </ol>	Calves	39,153
Cernivet LBC (Cerbios)	<i>Enterococcus faecium</i>	2 × 10 <sup>10</sup> cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased growth rate</li> <li>2. Increased forage and total dry matter intake</li> <li>3. Increased average weight gain</li> <li>4. Increased daily weight gain</li> </ol>	Lithuanian Black-and-White calves	132
Cylactin (DSM)	<i>E. faecium</i>	70 g kg <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved feed efficiency during pre-weaning</li> <li>2. Increased average daily gain</li> <li>3. Greater intake of crude protein and neutral detergent fiber</li> </ol>	Holstein calves	154
Protexin® (Protexin)	<i>Lactobacillus plantarum</i> , <i>L. bulgaricus</i> , <i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>Bifidobacterium bifidum</i> , <i>Streptococcus thermophilus</i> , <i>E. faecium</i> , <i>Aspergillus oryza</i> and <i>Candida pintolopesii</i>	2 × 10 <sup>8</sup> cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increase in average daily gain</li> <li>2. Increase in feed intake</li> <li>3. Increase LAB count</li> </ol>	Holstein female calves	155
Oralin® (Chevita GmbH)	<i>E. faecium</i> DSM 10663	1.5 × 10 <sup>7</sup> cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Control dysbiosis and intestinal disturbances</li> <li>2. Reconstituted the gut flora after antibiotic therapy</li> </ol>	Calves	2
PrimaLac (Star Labs, Inc.)	<i>B. bifidum</i> , <i>B. thermophilus</i> , <i>E. faecium</i> , <i>L. acidophilus</i> , <i>L. casei</i> ,	2.5 × 10 <sup>7</sup> cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Higher milk yield</li> <li>2. Increased milk components (lactose, protein and fat)</li> <li>3. Increased aspartate aminotransferase activity in blood plasma</li> </ol>	Cattle (beef, dairy), deer, ewe	156
Biovet –YC Gold (Biovet Health Products)	<i>L. acidophilus</i> , <i>Saccharomyces cerevisiae</i> , <i>S. boulardii</i> and <i>Propionibacterium freudenreichii</i>	20 g day <sup>-1</sup> cow <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased milk production of lactating cows</li> <li>2. Higher Milk fat, milk protein and Solids-Not Fat content</li> </ol>	Calves	33
No information provided	<i>L. casei</i>	1.0 × 10 <sup>7</sup> cfu g cow <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased the resistance of animals to adverse environmental conditions (high temperature and humidity)</li> <li>2. Increased milk yield in summer and winter</li> </ol>	Holstein-Friesian milk cow	157

**Table 1.** Continued

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
Pro-Biotyk EM15® (ProBiotics)	<i>B. subtilis</i> , <i>B. animalis</i> , <i>B. bifidum</i> , <i>B. longum</i> , <i>L. acidophilus</i> , <i>L. casei</i> , <i>delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. fermentum</i> , <i>L. plantarum</i> , <i>L. lactis</i> subsp. <i>Lactis</i> , <i>S. cerevisiae</i> , <i>S. thermophilus</i>	20 cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved feed digestibility</li> <li>2. Decreased emission of gases such as ammonia, hydrogen sulphide and mercaptans.</li> <li>3. Improved balance of the microflora in the digestive tract,</li> </ol>	Cattel, lambs, Calves	158
Pro-Biotyk EM15® (ProBiotics)	<i>B. subtilis</i> , <i>B. animalis</i> , <i>B. bifidum</i> , <i>B. longum</i> , <i>L. acidophilus</i> , <i>L. casei</i> , <i>L. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. fermentum</i> , <i>L. plantarum</i> , <i>L. lactis</i> subsp. <i>lactis</i> , <i>S. cerevisiae</i> , <i>S. thermophilus</i>	20 mL L <sup>-1</sup> water	<ol style="list-style-type: none"> <li>1. Improved balance of the microflora in the digestive tract,</li> <li>2. Reduction and stabilization of somatic cells in milk.</li> <li>3. Improved immunity</li> </ol>	Cattel, lambs, Calves	158
Yea Sacc (Alltech)	<i>S. cerevisiae</i>	1 × 10 <sup>9</sup> cfu cow <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved milk production and milk quality</li> <li>2. Increased energy corrected milk production</li> <li>3. increased synthesis of milk protein, milk protein content, and milk urea content</li> <li>4. Lower milk fat content</li> <li>5. Higher lactose content</li> <li>6. Enhanced energy utilization</li> <li>7. Reduced methane emissions</li> </ol>	lactating dairy cows	159
Yea-Sacc-1026 (Alltech)	<i>S. cerevisiae</i>	1 × 10 <sup>9</sup> cfu head <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved milk quality (fat and protein content)</li> <li>2. Increased body weight of lambs</li> </ol>	Lambs	73
Yea-Sacc-1026 (Alltech)	<i>S. cerevisiae</i>	2.5 × 10 <sup>9</sup> cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved feed efficiency</li> <li>2. Increased dry matter digestibility</li> </ol>	Dairy Holstein heifers	168
Yea-Sacc-1026 (Alltech)	<i>S. cerevisiae</i>	4.5 × 10 <sup>9</sup> cfu	<ol style="list-style-type: none"> <li>1. Increased in dry matter intake</li> <li>2. Increased digestion of non-digestible fiber</li> </ol>	Holstein heifers	168
Yea-Sacc-1026 (Alltech)	<i>S. cerevisiae</i>	2.5 × 10 <sup>9</sup> cfu g <sup>-1</sup> yeast	And hemicellulose digestibility	Multiparous dairy cows	169
No information provided	<i>L. acidophilus</i> , <i>L. salivarius</i> , <i>L. reuteri</i>	2.3 × 10 <sup>7</sup> cfu cow <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved metabolic activity</li> <li>2. Increased body weight</li> </ol>	Maltese goat kids	143
No information provided	<i>B. licheniformis</i> , <i>B. subtilis</i>	1 × 10 <sup>7</sup> cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased milk yield</li> <li>2. Higher milk quality</li> <li>3. Increased body weight</li> </ol>	Lambs	116

Table 1. Continued

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
No information provided	<i>Aspergillus oryzae</i>	$10^9$ cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. increased production of milk</li> <li>2. Increased ratio of protein and SNF in milk</li> </ol>	Cows	110
No information provided	<i>A. oryzae</i>	$5 \times 10^7$ cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Higher total volatile fatty acids, propionate, and acetate concentrations in the rumen.</li> <li>2. Higher counts of cellulolytic bacteria</li> </ol>	Calves	160
No information provided	<i>L. acidophilus</i>	$5 \times 10^7$ cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased body weight</li> </ol>	Calves	161
No information provided	<i>B. pseudolongum</i> , <i>L. acidophilus</i>	$3 \times 10^9$ cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved average daily gain</li> <li>2. Improved feed efficiency</li> </ol>	Calves	162
No information provided	<i>L. acidophilus</i> , <i>S. cerevisiae</i>	$1 \times 10^9$	<ol style="list-style-type: none"> <li>1. Increased average daily gain</li> </ol>	Calves	163
No information provided	<i>L. casei</i> DSPV 318 T, <i>L. salivarius</i> DSPV 315 T, <i>Pediococcus acidilactici</i> DSPV 006 T	$3 \times 10^9$ cfu flask <sup>-1</sup> kg <sup>-1</sup> $3 \times 10^9$ cfu kg <sup>-1</sup> live weight	<ol style="list-style-type: none"> <li>1. Stimulated consumption of Starter</li> <li>2. Early development of the rumen.</li> </ol>	Calves	68
No information provided	<i>E. faecium</i> , <i>L. plantarum</i> , <i>S. cerevisiae</i>	from $1 \times 10^5$ cfu mL <sup>-1</sup> to $1 \times 10^7$ cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Improved digestion rate</li> <li>2. Increased digestion of high moisture ear corn (HMEC) dry matter</li> </ol>	Cows	54
No information provided	<i>Propionibacterium</i> P15, <i>E. faecium</i> EF212	$1 \times 10^9$ cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>3. Reduction in intestinal pH</li> <li>1. Improved intake of dry matter</li> <li>2. Lowered blood CO<sub>2</sub> concentrations</li> </ol>	Cows	61
No information provided	<i>E. faecium</i> , Yeast	$5 \times 10^9$ cfu g <sup>-1</sup> $5 \times 10^9$ cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased consumption of dry matter</li> </ol>	Cows	164
No information provided	<i>Propionibacterium</i> P169	$6 \times 10^{10}$ cfu cow <sup>-1</sup> $6 \times 10^{11}$ cfu cow <sup>-1</sup>	<ol style="list-style-type: none"> <li>2. Increased daily milk production</li> <li>1. Increased milk yield</li> <li>2. Improved feed efficiency</li> <li>2. Increase daily gain</li> </ol>	Cows	57
No information provided	<i>L. acidophilus</i> LA747, <i>P. freudenreichii</i> PF24, <i>L. acidophilus</i> LA45	$1 \times 10^9$ cfu cow <sup>-1</sup> $2 \times 10^9$ cfu cow <sup>-1</sup> $5 \times 10^8$ cfu cow <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased milk yield</li> <li>2. increased fat corrected milk</li> <li>3. Reduced ruminal pH and total VFA concentration in the rumen</li> </ol>	Cows	165
No information provided	<i>E. faecium</i> , <i>S. cerevisiae</i>	$5 \times 10^9$ cfu cow <sup>-1</sup> day <sup>-1</sup> $5 \times 10^9$ cfu cow day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased in milk fat</li> <li>2. Received fewer antibiotic treatments.</li> </ol>	Cows	28
No information provided	<i>S. cerevisiae</i> subspecies <i>boulardii</i> CNCM I-1079	0.5 g of yeast steer <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Treatments did not affect dry matter intake, average daily gain or feed efficiency during the experimental period.</li> </ol>	Beef cattle	166



**Table 1.** Continued

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
No information provided	<i>Prevotella bryantii</i>	$2 \times 10^{11}$ cfu dose <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased milk fat</li> <li>2. Higher concentration of acetate, Butyrate</li> <li>3. Decreased lactate concentration</li> </ol>	Dairy cow	62
No information provided	<i>Propionibacterium</i> strain P169	$6 \times 10^{11}$ cfu day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Lowered concentrations of acetate</li> <li>2. Greater concentrations of propionate</li> <li>3. Higher energetic efficiency</li> </ol>	Cows	58
No information provided	<i>Propionibacterium</i> strain P169, yeast culture	$6 \times 10^{11}$ cfu steer <sup>-1</sup> day <sup>-1</sup> 56 g steer <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased molar proportions of propionate</li> <li>2. No effect on ruminal digestibility, microbial N synthesis, or particulate passage rates.</li> </ol>	Beef cattle	167
No information provided	<i>B. licheniformis</i> , <i>B. subtilis</i>	$2 \times 10^{11}$ cfu dose <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Reduction in mortality</li> <li>2. Increased milk production</li> </ol>	Sheep and Lambs	116
No information provided	<i>B. licheniformis</i> , <i>B. subtilis</i>	$2 \times 10^9$ cfu cow <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased milk yield and milk protein</li> <li>2. Increased ruminal digestibility</li> <li>3. Increased total volatile fatty acids concentration</li> </ol>	Holstein cows	111
No information provided	<i>B. licheniformis</i> , <i>B. subtilis</i>	$2 \times 10^9$ cfu head <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. higher average daily gain</li> <li>2. Increased final live weight</li> </ol>	Holstein Calves	153
No information provided	<i>S. cerevisiae</i> , QAU03, (locally isolated yeast from Sahiwal cow dung sample)	$3.13 \times 10^7$ cfu g <sup>-1</sup> yeast	<ol style="list-style-type: none"> <li>1. Increased fiber digestibility</li> <li>2. Improved milk and its fat contents</li> <li>3. Improved gastrointestinal tract microbial balance</li> </ol>	Lactating cows	168
Levucell 5c (Pasteur institute)	<i>S. cerevisiae</i> CNCM I-1077	$1 \times 10^{10}$ cfu head <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Enhanced rumen fermentation</li> <li>2. Lowered CH4 emissions</li> </ol>	Non-lactating dairy Holstein cows	87
Levucell Sc 20 (Pasteur institute)	Dry yeast (CNCM-1077)	0.5 g head <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Decreased time spent in subacute rumen acidosis</li> </ol>	Holstein dairy cows in late lactation	25
Levucell SC 10 ME (Pasteur institute)	<i>S. cerevisiae</i> CNCM I-1077	$1 \times 10^{10}$ cfu g <sup>-1</sup> yeast	<ol style="list-style-type: none"> <li>1. Lowered the risk of rumen acidosis</li> <li>2. Increased fiber</li> </ol>	Holstein dairy cows in Non-lactation	97

**Table 1.** Continued

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
Diamond-V XP yeast culture (Diamond-V)	<i>S. cerevisiae</i>	57 g day <sup>-1</sup> head <sup>-1</sup>	<p>degradation of low-quality maize silages</p> <ol style="list-style-type: none"> <li>1. Increased dry matter and protein digestion</li> <li>2. Increased propionate production</li> <li>3. Decreased microbial protein</li> </ol>	Lactating cows	170
A-Max yeast culture concentrate (A-Max)	<i>S. cerevisiae</i>	57 g day <sup>-1</sup> head <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased dry matter protein digestion</li> <li>2. Increased propionate percentage</li> <li>3. Decreased rumen pH</li> <li>4. Increased microbial nitrogen digestion</li> </ol>	Lactating cows	170
BIOSAF SC 47 (Lesaffre Feed Additives)	<i>S. cerevisiae</i> Sc47	5 g day <sup>-1</sup> head <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased volatile fatty acids concentrations</li> <li>2. Reduction rumen pH</li> <li>3. Increased counts of fibrolytic and lactate-using bacteria</li> </ol>	Early lactating cows	171
Actisaf Sc 47 (Phileo by Lesaffre)	<i>S. cerevisiae</i>	10 g day <sup>-1</sup> head <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Decreased acetate: propionate ratio</li> <li>2. Decreased Bacteroidetes</li> <li>3. Increased Fibrobacter bacteria</li> </ol>	Mid-to-late lactating cows	172
No information provided	<i>Orpinomyces</i> strain KNGF-2	200 mL head <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Increased nutrient digestibility and nitrogen retention</li> <li>2. Increased bacteria and fungi abundance</li> </ol>	Mature sheep	173
No information provided	<i>Orpinomyces</i> spp. C-14, <i>Piromyces</i> spp. WNG-12	160 mL day <sup>-1</sup> head <sup>-1</sup> for buffalo calves 250 mL week <sup>-1</sup> head <sup>-1</sup> for lactating buffaloes	<ol style="list-style-type: none"> <li>1. Increased digestibility of dry matter, crude protein, non-digestible fibre and cellulose</li> <li>2. Increased production of volatile fatty acids total nitrogen and trichloroacid precipitable nitrogen,</li> <li>3. Decreased ruminal pH and ammonia nitrogen</li> </ol>	Buffalo calves and lactating buffaloes	174,175
MTB-100 (Alltech)	Yeast-derived cell wall preparation	20, 40, or 60 g day <sup>-1</sup> head <sup>-1</sup>	<ol style="list-style-type: none"> <li>3. Increased zoospores</li> <li>1. Decreased ruminal indigestible</li> <li>2. Decreased fescue toxicosis</li> </ol>	Beef steers and cows	176
Solid powder (NA)	<i>S. cerevisiae</i> fermentation product	6 g kg <sup>-1</sup> DMI	<ol style="list-style-type: none"> <li>1. Decreased ruminal ammonia nitrogen</li> </ol>	Growing goats	177
No information provided	<i>L. casei</i> ssp. <i>casei</i>	1 × 10 <sup>8</sup> cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>2. Decreased methane emissions</li> <li>1. Increased weight gain</li> </ol>	Calves	98

**Table 1.** Continued

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
No information provided	Lactate-utilizing/or lactate-producing bacteria	1.3x10 <sup>6</sup> cfu mL <sup>-1</sup>	2. Improved feed efficiency 1. Improved feed efficiency 2. Increased in daily gain	Cattle	23
No information provided	Live yeast	60 g day <sup>-1</sup> head <sup>-1</sup>	1. Improved average daily gain 2. Increased final weight 3. Increased feed intake	Beef cattle	102
No information provided	Yeast	10 g day <sup>-1</sup> head <sup>-1</sup>	3. Improved feed to gain ratio 1. Increased milk yield and milk quality 2. Increased feed efficiency 3. Reduced ruminal acidosis	Dairy cows	57,100
No information provided	<i>L. casei</i> Zhang and <i>L. plantarum</i> P-8	1 × 10 <sup>8</sup> cfu mL <sup>-1</sup>	1. Improved quality and quantity of milk production Improve body weight	Dairy cows	109
FasTrack Microbial pack (Conklin fastrack)	<i>L. acidophilus</i> , <i>S. cerevisiae</i> , <i>E. faecium</i> , <i>A. oryzae</i> , active dry yeast culture	5 g day <sup>-1</sup> head <sup>-1</sup>		Dairy cows	44,96
No information provided	<i>L. acidophilus</i> , <i>S. cerevisiae</i> , <i>S. boulardii</i> , <i>P. freudenreichii</i>	5 g day <sup>-1</sup> head <sup>-1</sup>	1. Increased milk production 2. Increased feed intake	Lactating cows	33
No information provided	<i>B. licheniformis</i> and <i>B. subtilis</i>	1 × 10 <sup>8</sup> cfu mL <sup>-1</sup>	1. Reduced lamb mortality rate 2. Increased average daily 3. Milk yield per ewe 4. Improved fat and protein content of milk	Sheep	116
No information provided	<i>S. cerevisiae</i>	2.5 × 10 <sup>6</sup> cfu mL <sup>-1</sup>	1. Increased in the mean protozoal count 2. Increased total bacterial count	Buffalo bulls	178
No information provided	Live Bacillus culture: <i>B. licheniformis</i>	100 g day <sup>-1</sup>	1. Increased in microbial crude protein flow into duodenum 2. Decreased in ammonia nitrogen concentration in the ruminal fluid 3. Increased total volatile fatty acids and acetate concentrations in the ruminal fluid	Chinese Holstein cows	111

Table 1. Continued

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
No information provided	<i>Ruminococcus flavefaciens</i> NJ	$1 \times 10^6$ cfu mL <sup>-1</sup>	<p>4. Improved ruminal apparent nutrient digestibility of non-digestible fats</p> <p>1. Increased abundance of cellulolytic bacterial populations in the rumen</p> <p>2. Improved in sacco digestibility of Timothy hay with the high concentrate diet.</p> <p>3. Increased in the persistence of beneficial bacteria <i>R. flavefaciens</i> NJ</p>	Dairy cows	62
No information provided	A mixture of <i>L. plantarum</i> , <i>P. acidilactici</i> , <i>E. faecium</i> and <i>L. lactis</i>	$5 \times 10^5$ cfu g <sup>-1</sup> fresh herbage	<p>1. Increased in lactic acid concentration</p> <p>2. Decreased concentration of acetic acid.</p> <p>2. Higher rumen microbe count</p> <p>3. Lowered ratio of acetic acid to propionic acid.</p> <p>4. Improved rumen protein synthesis</p> <p>5. Higher content of protein nitrogen and</p>	Dairy cows	179
No information provided	<i>P. acidilactici</i> and <i>P. pentosaceus</i>	$1 \times 10^8$ cfu mL <sup>-1</sup>	<p>total nitrogen in the rumen.</p> <p>1. Improved average daily gain</p> <p>2. Increased final body weight</p> <p>3. Improved feed conversion ratio</p>	Lambs during post-weaning period	101
No information provided	<i>L. acidophilus</i>	$10^6$ cfu mL <sup>-1</sup>	<p>Increased the average daily weight gain</p>	buffalo calves	104
No information provided	<i>S. cerevisiae</i>	$4 \times 10^9$ cfu day <sup>-1</sup> animal <sup>-1</sup>	<p>Increased in the average daily milk yield</p>	lactating Saanen dairy goats	113
No information provided	<i>L. acidophilus</i> and <i>A. awamori</i>	$10^8$ cfu mL <sup>-1</sup>	<p>Improved digestibility of organic matter, dry matter, crude fiber, crude protein and nitrogen-free extract</p>	Goats	126
No information provided	<i>L. acidophilus</i> and <i>P. freudenreichii</i>	$2.5 \times 10^4$ cfu mL <sup>-1</sup>	<p>Decreased incidences of diarrhoea and other diseases of GIT</p>	lambs and calves	64

**Table 1.** Continued

Probiotic product (manufacturer)	Probiotic strain(s)	Probiotic dosage	Impacts	Animal species (host)	Reference
CSPB (calf specific probiotic) (Winclove)	<i>L. acidophilus</i> W55, <i>L. salivarius</i> W57, <i>L. paracasei</i> spp. <i>paracasei</i> W56, <i>L. plantarum</i> W59, <i>L. lactis</i> W58, and <i>E. faecium</i> W54 <i>Lactobacillus</i> spp.	$1.0 \times 10^9$ cfu kg <sup>-1</sup>	<ol style="list-style-type: none"> <li>Enhanced growth rate</li> <li>Increased body weight gain</li> <li>Improved feed consumption</li> <li>Decrease mortality</li> </ol>	Veal calves	180
Lactoamylorinum (NA)	<i>L. acidophilus</i> spp.	1 mL day <sup>-1</sup> head <sup>-1</sup>	<ol style="list-style-type: none"> <li>Increased daily weight gain</li> <li>Improved nutrient digestibility</li> <li>Improved carcass quality</li> <li>Decreased cost per unit gain in live weight</li> </ol>	Calves	181
No information provided	<i>L. acidophilus</i> , <i>L. jugarti</i> , <i>L. casei</i>	$1.0 \times 10^8$ cfu kg <sup>-1</sup>	<ol style="list-style-type: none"> <li>Increased daily weight gain</li> <li>Improved feed conversion and lower feed intake</li> <li>Improved daily weight gain</li> <li>Significantly increased milk production</li> <li>Increased the concentration of acetic acid</li> <li>Improved the aerobic stability of the ration in lactating cows</li> </ol>	Calves	41,182
No information provided	<i>L. acidophilus</i>	$2.5 \times 10^7$ cfu	<ol style="list-style-type: none"> <li>Improved daily weight gain</li> <li>Improved feed conversion</li> </ol>	Holstein calves	183
No information provided	<i>L. acidophilus</i> , <i>L. buchneri</i> 40 788	$1 \times 10^5$ , $5 \times 10^5$ , or $1 \times 10^6$ cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>Significantly increased milk production</li> <li>Increased the concentration of acetic acid</li> <li>Improved the aerobic stability of the ration in lactating cows</li> <li>Increased milk production</li> </ol>	Dairy cows, lactating cows	55
No information provided	<i>L. acidophilus</i>	$10^7$ cfu g <sup>-1</sup>	<ol style="list-style-type: none"> <li>Increased daily milk yields</li> <li>Increased milk protein content</li> </ol>	Cows	184
No information provided	<i>L. acidophilus</i> , <i>S. faecium</i> , <i>L. casei</i> , <i>L. fermentum</i> and <i>L. plantarum</i>	$10^{10}$ cfu	<ol style="list-style-type: none"> <li>Stimulated feed intake</li> <li>Improved daily weight gain</li> </ol>	Lambs	185
NA, Not available.					

**Table 2.** Summary of probiotics effects on ruminant health

Probiotic product (Manufacturer)	Probiotic strain(s)	Probiotic dose	Impacts	Ruminant host	Reference
No information provided	<i>Lactobacillus acidophilus</i> , <i>Propionibacterium freudenreichii</i>	From $1 \times 10^6$ cfu mL <sup>-1</sup> to $1 \times 10^9$ cfu mL <sup>-1</sup>	Lowered faecal shedding of <i>E. coli</i>	Calves	186
No information provided	<i>Bacillus cereus</i> var. Toyoi <i>Saccharomyces boulardii</i>	$1.0 \times 10^8$ cfu kg <sup>-1</sup>	Both probiotics enhanced humoral immunity.	Sheep	187
No information provided	<i>Bacillus subtilis</i> strain 166	$1 \times 10^9$ cfu mL <sup>-1</sup>	1. There were no significant differences observed between treatments for either hide or fecal prevalence of <i>E. coli</i> O157: H7.	Cattle	188
Calf-specific probiotic (NA)	<i>L. acidophilus</i> W55, <i>L. salivarius</i> W57, <i>L. paracasei</i> spp. <i>paracasei</i> W56, <i>L. plantarum</i> W59, <i>Lactococcus</i> <i>lactis</i> W58, and <i>Enterococcus</i> <i>faecium</i> W54.	$1.0 \times 10^9$ cfu kg <sup>-1</sup>	1. Reduced diarrhea 2. Decreased fecal coliform counts	Veal calves	180
No information provided	<i>L. acidophilus</i> , <i>S. cerevisiae</i> , <i>E. faecium</i> , <i>Aspergillus oryzae</i> , <i>fructooligosaccharide</i> , active dry yeast culture	$2.50 \times 10^8$ cfu g <sup>-1</sup>	1. Improved Packed cell volume 2. Increased Faffa Malan Chart (FAMACHA) scorescores	Goats	50
No information provided	<i>L. acidophilus</i> and <i>P. freudenreichii</i>	$1 \times 10^9$ cfu mL <sup>-1</sup>	1. Decreased incidences of diarrhoea 2. Decreased diseases of GIT	Lambs and calves	64
Cernivet LBC (Cerbios)	<i>E. faecium</i>	$2.4 \text{ g day}^{-1} \text{ calve}^{-1}$	1. Reduction in diarrhoea 2. Reduction in faecal count of clostridia and enterococci.	Calves	132
No information provided	<i>L. plantarum</i> , <i>bulgaricus</i> , <i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>Bifidobacterium bifidum</i> , <i>Streptococcus thermophilus</i> , <i>E. faecium</i> , <i>Aspergillus oryza</i> and <i>Candida pintolopesii</i>	$3.08 \times 10^8$ cfu g <sup>-1</sup>	1. Controlled diarrhoea and bloat 2. Reduction of mortality 3. Reduction in veterinary cost of treatment	Holstein male calves	189
Protexin® (Protexin)	<i>L. plantarum</i> , <i>L. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>B. bifidum</i> , <i>S. salivarius</i> subsp. <i>thermophilus</i> , <i>E. faecium</i> and yeast	$2 \times 10^8$ cfu g <sup>-1</sup>	1. Significantly lowered count of pathogenic <i>E. coli</i> 2. Mitigation of diarrhoea	Holstein female calves	155
Protexin (Protexin)	<i>L. plantarum</i> , <i>delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. acidophilus</i> , <i>L. Rhamnosus</i> , <i>Bifidobacterium</i> <i>bifidum</i> , <i>S. salivarius</i> subsp. <i>thermophilus</i> , <i>E. faecium</i> , <i>A. oryzae</i> , <i>C. pintolopesii</i>	$2 \text{ g day}^{-1} \text{ head}^{-1}$	Significant decreased in ileocecal infection by <i>Mycobacterium avium</i> ssp. <i>paratuberculosis</i> (MAP)	Holstein bull calves	190

**Table 2.** Continued

Probiotic product (Manufacturer)	Probiotic strain(s)	Probiotic dose	Impacts	Ruminant host	Reference
No information provided	<i>L. casei</i>	$1.0 \times 10^7$ cfu <sup>-1</sup> g cow <sup>-1</sup> day <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Decreased in the number of somatic cells</li> <li>2. Decreased in the incidence of mastitis</li> <li>3. Positive effect on the microflora of the gastrointestinal tract</li> <li>3. Greater resistance to infectious diseases</li> </ol>	Holstein-Friesian milk cow	157
No information provided	<i>L. acidophilus</i>	$10^9$ cfu steer <sup>-1</sup> daily	Reduced <i>E. coli</i> O157 levels	Beef cattle, steer	131
No information provided	<i>L. acidophilus</i> strain NP 51	$10^9$ cfu steer <sup>-1</sup> daily	Reduced <i>E. coli</i> O157 prevalence in both fecal and hide sample	Beef cattle	191
CSPB (calf specific probiotic) (Winclove)	<i>L. acidophilus</i> W55, <i>L. salivarius</i> W57, <i>L. paracasei</i> spp. <i>paracasei</i> W56, <i>L. plantarum</i> W59, <i>L. lactis</i> W58, and <i>E. faecium</i> W54	$1.0 \times 10^9$ cfu kg <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Decreased mortality</li> <li>2. Reduced the incidence of diarrhea</li> <li>3. Reduced faecal counts of coliforms</li> <li>4. Reduction in the percentage of calves in need of therapy and the amount of treatments needed against digestive or respiratory diseases</li> </ol>	Veal calves	180
No information provided	<i>L. acidophilus</i> NPC 747	$1 \times 10^9$ cfu	<ol style="list-style-type: none"> <li>1. Decreased but did not eliminated fecal shedding of <i>E. coli</i> O157:H7</li> <li>2. Reduced <i>E. coli</i> O157:H7 contamination on hides</li> </ol>	Cattle	192
No information provided	<i>L. gallinarum</i> LCB 12, <i>S. bovis</i>	$1 \times 10^{10}$ cfu	<ol style="list-style-type: none"> <li>1. Reduced or stopped carriage of <i>E. coli</i> O157</li> </ol>	Holstein calves	193,194
No information provided	<i>L. acidophilus</i>	$2.5 \times 10^7$ cfu	Decreased diarrhea	Holstein calves	183
No information provided	<i>L. acidophilus</i> , <i>S. faecium</i> , <i>L. casei</i> , <i>L. fermentum</i> and <i>L. plantarum</i>	$10^{10}$ cfu	Reduced fecal shedding of <i>E. coli</i> O157:H7	Lambs	185
No information provided	<i>B. pseudolongum</i> , <i>L. acidophilus</i>	$3 \times 10^9$ cfu mL <sup>-1</sup>	Reduction in diarrhea incidence	Calves	162
No information provided	<i>L. acidophilus</i> , <i>L. plantarum</i> , <i>L. acidophilus</i> 275C	$1.85 \times 10^7$ cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Decreased incidence of diarrhea</li> <li>2. Increased lactobacilli count in feces.</li> </ol>	Calves	195
Oralin® (Chevita GmbH)	<i>E. faecium</i> DSM 10663	$2.5 \times 10^8$ cfu mL <sup>-1</sup>	<ol style="list-style-type: none"> <li>1. Controlled dysbiosis and intestinal disturbances</li> <li>2. Reconstituted the gut flora after antibiotic therapy</li> </ol>	Calves	2

**Table 2.** Continued

Probiotic product (Manufacturer)	Probiotic strain(s)	Probiotic dose	Impacts	Ruminant host	Reference
Pro-Biotyk EM15® (ProBioTics)	<i>B. subtilis</i> , <i>B. animalis</i> , <i>B. bifidum</i> , <i>B. longum</i> , <i>L. acidophilus</i> , <i>L. casei</i> , <i>L. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. fermentum</i> , <i>L. plantarum</i> , <i>L. lactis</i> subsp. <i>lactis</i> , <i>S. Cerevisiae</i> , <i>S. thermophilus</i>	20 mL L <sup>-1</sup> water	1. Improved balance of the microflora in the digestive tract, 2. Reduction of diarrhea occurrence 3. Reduction and stabilization of somatic cells in milk.	Cattel, lambs, Calves	158
No information provided	<i>B. pseudolongum</i> , <i>L. acidophilus</i>	3 × 10 <sup>9</sup> cfu mL <sup>-1</sup>	3. Improved immunity and	Calves	162
No information provided	<i>Propionibacterium</i> P15, <i>E. faecium</i> EF212	1 × 10 <sup>9</sup> cfu g <sup>-1</sup>	1. Reduced diarrhea incidence. 1. Lowered blood CO <sub>2</sub> concentrations 3. Reduced risk of metabolic acidosis.	Cow	63
No information provided	<i>B. subtilis</i>	10 <sup>8</sup> cfu head <sup>-1</sup>	Reduced <i>Clostridium perfringens</i> shedding	Holstein calves	(Wöhnes <i>et al.</i> , 196)
No information provided	<i>L. acidophilus</i> and <i>P. freudenreichii</i>	1 × 10 <sup>6</sup> cfu mL <sup>-1</sup>	1. Decreased incidences of diarrhea 2. Decreased in the occurrence of diseases of gastrointestinal tract	Lambs and calves	64
NA, Not available.					

frequency, animal species, animal age and physiological state, environmental conditions, etc., can influence the effectiveness of probiotics in the improvement of nutrient digestibility and feed intake in ruminants<sup>2,127</sup> (Table 1).

#### Pathogen reduction and gastrointestinal disease mitigation

Apart from the fact that some ruminants are the primary reservoir host for some foodborne pathogens, outbreaks of some human diseases are linked to some ruminant species.<sup>1,120,128</sup> The transmission of these zoonotic pathogens from ruminants to humans constitute a serious public health problem with huge global economic loss.<sup>1,128</sup> In the last decade, scientists have increasingly sought out healthy and safe agricultural practices in livestock production, aimed at reducing the carriage of gut pathogens, their release unto the environment and their transmission to humans. Major zoonotic pathogens that have been targeted are *E. coli*, *E. coli* O157, *Campylobacter*, *Enterococcus*, *Listeria*, *Salmonella* and *Shigella* spp.<sup>25,128</sup> The dietary inclusion of probiotics has been shown to reduce the enteric colonization of pathogens in ruminants, which enhance overall animal health, as well as prevent human transmission and environmental contamination<sup>2,30,129,130</sup> (Table 2).

LAB, especially strains of *L. acidophilus*, used in feedlot cattle have been shown to significantly reduce the carriage and shedding of *E. coli* O157 and *Salmonella enterica*.<sup>131</sup> Shedding of *E. coli* O157:H7 was reduced in cattle after the dietary inclusion of a multi-strain probiotic containing *L. acidophilus* and *P. freudenreichii*.<sup>129</sup> Strains of LAB are most effective in reducing the carriage of gut pathogens in ruminants.<sup>128</sup> Incidences of diarrhoea and other diseases of the GIT that result in weight loss in young ruminants (especially lambs and calves) have been shown to subside after the application of LAB.<sup>30,64,132</sup> Stress in ruminants may cause an imbalance in the gut microflora or dysbiosis, which may be controlled through the administration of probiotics.<sup>2</sup> The application of dietary probiotics in livestock production not only reduces pathogen carriage and gut disease, but also further limits the risk of human infections with foodborne pathogens.<sup>130</sup>

#### Haemato-biochemical parameters and metabolites

There are conflicting reports on the effects of probiotics on haemato-biochemical parameters and metabolites levels in ruminants. Although some studies show no difference in glucose concentration in lambs supplemented with probiotic,<sup>51,133</sup> Antunović *et al.*<sup>134</sup> and Bruno *et al.*<sup>135</sup> reported significant decreases in glucose concentrations. Conversely, probiotic supplementation of lactating ewes and lambs significantly increased the concentration of glucose.<sup>126,136</sup> It was reported previously that the relationship existing between blood glucose levels and probiotic inclusion is dose-dependent.<sup>137</sup> Cetin *et al.*<sup>138</sup> observed a statistical increase in erythrocyte sedimentation rate, haemoglobin concentration and haematocrit values in animals supplemented with probiotics. The supplementation of either single strains of *L. lactis* and *L. plantarum* or their combination as multi-strain probiotics significantly increased total red blood cell counts and haemoglobin concentration in the blood.<sup>139</sup> The dietary inclusion of probiotics positively influenced haematopoiesis, which, amongst others, benefits increases in total white blood cell counts, hence enhancing immune cells synthesis, further protecting the host against invading pathogens.<sup>1,140</sup>

The findings from several studies revealed that supplementation with probiotics in animals may significantly improve the lipid profile in the blood of ruminants.<sup>141</sup> The supplementation of kids



or lambs with probiotics was shown to decrease the concentrations of triglycerides, total lipids, low-density lipoproteins and non-esterified fatty acids in the blood of ruminants.<sup>142,143</sup> Panda *et al.*<sup>144</sup> reported a significant reduction in total cholesterol and triglycerides by dietary inclusion of 100 mg kg<sup>-1</sup> diet of *Lactobacillus sporogenes* probiotic in animals. Total cholesterol reduction in probiotic supplemented animals could be the result of direct assimilation of cholesterol by bacterial cells (which causes a reduction in the cholesterol absorption and synthesis in the GIT), 3-hydroxy-3-methyl-glutaryl-CoA reductase inhibition and bile salt hydrolysis.<sup>145,146</sup> Furthermore, triglyceride reduction in probiotic treated animals may be a result of increased hydrolysis of bile salts, which causes inadequate lipid absorption in the small intestine.<sup>147</sup> Strains of *Lactobacillus* are known to show high hydrolytic activity of bile salts, which consequently leads to bile salt deconjugation within the GIT.<sup>148</sup> Several studies have reported a decrease in the concentration of certain blood metabolites including urea and blood urea nitrogen when lambs were fed with diets supplemented with probiotics.<sup>134,135</sup> Probiotic supplementation in lambs increases the bacterial population in the rumen, which may lead to improved utilization of nitrogen by ruminal bacteria thereby reducing blood urea nitrogen concentration<sup>135</sup> (Table 2).

## SUSTAINABILITY OF RUMINANT PRODUCTION

The importance of ruminants in the production of high volumes of meat and milk, which are important animal proteins for human nutrition and health, and their ability to effectively use fibrous non-digestible feed cannot be overemphasized.<sup>6</sup> Despite an increasing loss in soil fertility, desertification, erosion and drought, the vast majority of people in the developing world primarily depend on ruminants as their major food sources.<sup>149</sup> However, apart from this, the environmental impact of the emission of greenhouse gases by ruminants and other problems associated with their health and welfare constitutes a major global challenge.<sup>7</sup>

With the continued global rise in the human population, the supply of sufficient, high quality and safe food is threatened by a myriad of factors.<sup>150</sup> Food security and the sustainability of agriculture, especially livestock production, has become an immense concern for both developed and developing countries of the world.<sup>2</sup> Considering the role that ruminants play both in the production of human food and employment, there is an urgent need to chart a new course of action that will lead to a sustainable increase in the production of food of ruminant origin, as well as improve the health and welfare of animals.<sup>6,151</sup>

The increased interest in organic farming and consumers' preference for chemical and/or antibiotic-free animal products will lead to more sustainable and beneficial effects both in the livestock industry, as well as environmental and public health. Furthermore, the inclusion of probiotics and other natural, safe and novel feed sources in animal husbandry may lead to a significant improvement in the production of food which will also be more sustainable.<sup>152</sup> Although plant- and cell-based food are becoming more prevalent in human diets, ruminants are able to transform plant material that is undigestible by humans into animal foods of high nutritive value. If there is no global advancement for alternatives to the use of antibiotics in ruminant-based food production systems, negative impacts on their productivity could occur, leading to shortages in food production.<sup>149</sup> Advantages from

the use of probiotics could extend to systems that do not rely on the use of antibiotics, such as organic farming. Should the current trend in human population increase continue, by 2050, there will be less arable land, which will eventually create unhealthy competition, overexploitation and a decline in biodiversity. As such, an improvement and intensification of sustainable agriculture and livestock production will be the only realistic solution to meet human needs.<sup>149</sup> The global acceptance of sustainable ruminant production through supplementation with probiotics may thus help maintain food security and food safety for the world.

## FUTURE DIRECTION AND CONCLUSIONS

The supplementation of ruminants with probiotics can contribute immensely to improved growth, general performance and overall health status. However, continuous in-depth, comprehensive and encompassing *in vivo* studies that use high-throughput omic technologies to fully decipher the mechanisms and effects exerted by probiotics on ruminants and their role in improving animal health and nutrition, as well as the subsequent effect on the sustainability of food production, are needed. Also, the characterization of metabolites secreted by probiotic strains and their role in improving ruminants' health and nutrition should be explored.

## ACKNOWLEDGEMENTS

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## REFERENCES

- Gaggia F, Mattarelli P and Biavati B, Probiotics and prebiotics in animal feeding for safe food production. *Int J Food Microbiol* **141**:S15–S28 (2010). <https://doi.org/10.1016/j.jfoodmicro.2010.02.031>.
- FAO, *Probiotics in Animal Nutrition: Production, Impact and Regulation*. FAO, Rome (2016).
- Smith J, Sones K, Grace D, MacMillan S, Tarawali S and Herrero M, Beyond milk, meat, and eggs: role of livestock in food and nutrition security. *Anim Front* **3**:6–13 (2013). <https://doi.org/10.2527/af.2013-0002>.
- UN, 2018. United Nations (UN). 2019. Department of Economic and Social Affairs. Population Division. World Population Prospects, the 2019 revision [Internet].
- Herrero M, Wirsenius S, Henderson B, Rigolot C, Thornton P, Havlík P *et al.*, Livestock and the environment: what have we learned in the past decade? *Annu Rev Env Resour* **40**:177–202 (2015). <https://doi.org/10.1146/annurev-environ-031113-093503>.
- Eisler MC, Lee MRF, Tarlton JF and Martin GB, Steps to sustainable livestock. *Nature* **507**:32–34 (2014). <https://doi.org/10.1038/507032a>.
- Knapp JR, Laur GL, Vadas PA, Weiss WP and Tricarico JM, Invited review: enteric methane in dairy cattle production: quantifying the opportunities and impact of reducing emissions. *J Dairy Sci* **97**:3231–3261 (2014). <https://doi.org/10.3168/jds.2013-7234>.
- Ogbuewu IP, Okoro VM, Mbajiorgu EF and Mbajiorgu CA, Yeast (*Saccharomyces cerevisiae*) and its effect on production indices of livestock and poultry—a review. *Comp Clin Path* **28**:669–677 (2019). <https://doi.org/10.1007/s00580-018-2862-7>.
- Wegener HC, Antibiotics in animal feed and their role in resistance development. *Curr Opin Microbiol* **6**:439–445 (2003). <https://doi.org/10.1016/j.mib.2003.09.009>.
- Jouany JP and Morgavi DP, Use of “natural” products as alternatives to antibiotic feed additives in ruminant production. *Animal* **1**:1443–1466 (2007). <https://doi.org/10.1017/S1751731107000742>.
- Russell JB and Houlihan AJ, Ionophore resistance of ruminal bacteria and its potential impact on human health. *FEMS Microbiol Rev* **27**:65–74 (2003). [https://doi.org/10.1016/S0168-6445\(03\)00019-6](https://doi.org/10.1016/S0168-6445(03)00019-6).

- 12 Khan R, Ullah Khan R, Naz S, Dhama K, Karthik K, Tiwari R *et al.*, Direct-fed microbial: beneficial applications, modes of action and prospects as a safe tool for enhancing ruminant production and safeguarding health. *Int J Pharmacol* **12**:220–231 (2016). <https://doi.org/10.3923/ijp.2016.220.231>.
- 13 Ayala DI, Cook PW, Franco JG, Bugarel M, Kottapalli KR, Loneragan GH *et al.*, A systematic approach to identify and characterize the effectiveness and safety of novel probiotic strains to control foodborne pathogens. *Front Microbiol* **10**:1108 (2019). <https://doi.org/10.3389/fmicb.2019.01108>.
- 14 Chichlowski M, Croom WJ, Edens FW, McBride BW, Qiu R, Chiang CC *et al.*, Microarchitecture and spatial relationship between bacteria and ileal, cecal, and colonic epithelium in chicks fed a direct-fed microbial, primaLac, and salinomycin. *Poult Sci* **86**:1121–1132 (2007). <https://doi.org/10.1093/PS/86.6.1121>.
- 15 van Zyl WF, Deane SM and Dicks LMT, Bacteriocin production and adhesion properties as mechanisms for the anti-listerial activity of *Lactobacillus plantarum* 423 and *Enterococcus mundtii* ST4SA. *Benef Microbes* **10**:329–349 (2019). <https://doi.org/10.3920/BM2018.0141>.
- 16 Wilkins T and Sequoia J, Probiotics for gastrointestinal conditions: a summary of the evidence. *Am Fam Physician* **96**(3):170–178 (2017).
- 17 Kollath W, The increase of the diseases of civilization and their prevention. *Munch Med Wochenschr* **95**:1260–1262 (1953).
- 18 Lilly DM and Stillwell RH, Probiotics: growth-promoting factors produced by microorganisms. *Science* (80- ) **147**:747–748 (1965). <https://doi.org/10.1126/science.147.3659.747>.
- 19 Parker RB, Probiotics, the other half of the antibiotic story. *Anim Nutr Health* **29**:4–8 (1974).
- 20 Fuller R, Probiotics in man and animals. *J Appl Bacteriol* **66**:365–378 (1989). <https://doi.org/10.1111/j.1365-2672.1989.tb05105.x>.
- 21 Kmet V, Flint HJ and Wallace RJ, Probiotics and manipulation of rumen development and function. *Arch Tierernahr* **44**:1–10 (1993). <https://doi.org/10.1080/17450399309386053>.
- 22 Miles RD and Bootwala SM, Direct-fed microbials in animal production—avian—a review, in *Direct-Fed Microbial in Animal Production*. National Feed Ingredients Association, West Des Moines, IA, pp. 117–146 (1991). <https://doi.org/10.3382/ps.0750186>.
- 23 Krehbiel C, Rust S, Zhang G and Gilliland S, Bacterial direct-fed microbials in ruminant diets: performance response and mode of action. *J Anim Sci* **81**:E120–E132 (2003). [https://doi.org/10.2527/2003.8114\\_suppl\\_2E120x](https://doi.org/10.2527/2003.8114_suppl_2E120x).
- 24 Yang Y, Iji PA and Choct M, Dietary modulation of gut microflora in broiler chickens: a review of the role of six kinds of alternatives to in-feed antibiotics. *Worlds Poult Sci J* **65**:97–114 (2009). <https://doi.org/10.1017/S0043933909000087>.
- 25 Chaucheyras-Durand F and Durand H, Probiotics in animal nutrition and health. *Benef Microbes* **1**:3–9 (2010). <https://doi.org/10.3920/BM2008.1002>.
- 26 Elghandour MMY, Chagoyán JCV, Salem AZM, Kholif AE, Castañeda JSM, Camacho LM *et al.*, Effects of *Saccharomyces cerevisiae* at direct addition or pre-incubation on *in vitro* gas production kinetics and degradability of four fibrous feeds. *Ital J Anim Sci* **13**:3075 (2014). <https://doi.org/10.4081/ijas.2014.3075>.
- 27 O'Connor MH, Martin SA and Hill GM, Effects of *Saccharomyces cerevisiae* on *in vitro* mixed ruminal microorganism fermentation. *Prof Anim Sci* **18**:358–362 (2002). [https://doi.org/10.15232/S1080-7446\(15\)31546-1](https://doi.org/10.15232/S1080-7446(15)31546-1).
- 28 Oetzel GR, Emery KM, Kautz WP and Nocek JE, Direct-fed microbial supplementation and health and performance of pre- and postpartum dairy cattle: a field trial. *J Dairy Sci* **90**:2058–2068 (2007). <https://doi.org/10.3168/jds.2006-484>.
- 29 Yoon IK and Stern MD, Influence of direct-fed microbials on ruminal microbial fermentation and performance of ruminants - a review. *Asian-Australas J Anim Sci* **8**:533–555 (1995). <https://doi.org/10.5713/ajas.1995.553>.
- 30 Seo JK, Kim S-W, Kim MH, Upadhaya SD, Kam DK and Ha JK, Direct-fed microbials for ruminant animals. *Asian-Australas J Anim Sci* **23**:1657–1667 (2010). <https://doi.org/10.5713/ajas.2010.r.08>.
- 31 FAO, Guidelines for the evaluation of probiotics in food, in *Report of a Joint FAO/WHO Working Group on Drafting Guidelines for the evaluation of probiotics in food*. FAO, London, Ontario, Canada (2002). [https://www.who.int/foodsafety/fs\\_management/en/probiotic\\_guidelines.pdf?ua=1](https://www.who.int/foodsafety/fs_management/en/probiotic_guidelines.pdf?ua=1)
- 32 Ezema C, Probiotics in animal production: a review. *J Vet Med Anim Health* **5**:308–316 (2013). <https://doi.org/10.5897/JVMAH2013.0201>.
- 33 Vibhute VM, Shelke RR, Chavan SD and Nage SP, Effect of probiotics supplementation on the performance of lactating crossbred cows. *Vet World* **4**:557–561 (2011). <https://doi.org/10.5455/vetworld.2011.557-561>.
- 34 Kruijs W, Frič P, Pokrotnieks J, Lukáš M, Fixa B, Kaščák M *et al.*, Maintaining remission of ulcerative colitis with the probiotic *Escherichia coli* Nissle 1917 is as effective as with standard mesalazine. *Gut* **53**:1617–1623 (2004). <https://doi.org/10.1136/gut.2003.037747>.
- 35 Russell P, Physiological and functional properties of probiotics (2008) - by international dairy federation. *Int J Dairy Technol* **62**:290–291 (2009). <https://doi.org/10.1111/j.1471-0307.2009.00488.x>.
- 36 Saarela M, Mogensen G, Fondén R, Mättö J and Mattila-Sandholm T, Probiotic bacteria: safety, functional and technological properties. *J Biotechnol* **84**:197–215 (2000). [https://doi.org/10.1016/S0168-1656\(00\)00375-8](https://doi.org/10.1016/S0168-1656(00)00375-8).
- 37 Morelli L, *In vitro* assessment of probiotic bacteria: from survival to functionality. *Int Dairy J* **17**:1278–1283 (2007). <https://doi.org/10.1016/j.idairyj.2007.01.015>.
- 38 Shanahan F, Probiotics: a perspective on problems and pitfalls, in *Scandinavian J. Gastroenterol.* Taylor & Francis. **38**:34–36 (2003). <https://doi.org/10.1080/00855910310001476>.
- 39 EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), Rychen G, Aquilina G, Azimonti G, Bampidis V, Bastos ML *et al.*, Guidance on the characterisation of microorganisms used as feed additives or as production organisms. *EFSA J* **16**:5206 (2018). <https://doi.org/10.2903/j.efsa.2018.5206>.
- 40 Ravi N, Pravin DS, Khedkar CD and Ajay S, Selection criteria for probiotics: a review. *Int J Probiotics Prebiotics* **9**:1–6 (2014).
- 41 Dash SK, Selection criteria for probiotics, in *XXXVII Dairy Industry Conference, Kala Academy, Panjim, Goa, 1–4, Feb 7–9, 2009*, ed. by Dash SK. Kala Academy, Panjim (2009). XXXVII Dairy Industry Conference
- 42 Elmer GW, McFarland LV and McFarland M, The power of probiotics: Improving your health with beneficial microbes, in *The Power of Probiotics: Improving Your Health with Beneficial Microbes*. Taylor and Francis, New York, USA (2013). <https://doi.org/10.4324/9781315821139>.
- 43 Schrezenmeier J and de Vrese M, Probiotics, prebiotics, and synbiotics - approaching a definition. *Am J Clin Nutr* **73**:361s–364s (2001). <https://doi.org/10.1093/ajcn/73.2.361s>.
- 44 Adjei-Fremah S, Ekwemalor K, Asiamah E, Ismail H and Worku M, Transcriptional profiling of the effect of lipopolysaccharide (LPS) pre-treatment in blood from probiotics-treated dairy cows. *Genomics Data* **10**:15–18 (2016). <https://doi.org/10.1016/j.gdata.2016.08.016>.
- 45 Ekwemalor K, Adjei-Fremah S, Asiamah E, Ismail H and Worku M, 0167 exposure of bovine blood to pathogen associated and non pathogen associated molecular patterns results in transcriptional activation. *J Anim Sci* **94**:81–81 (2016). <https://doi.org/10.2527/jam2016-0167>.
- 46 Gyenai KB, Worku M, Tajkarimi M and Ibrahim S, Influence of probiotics on coccidia, *H. contortus* and markers of infection in goats. *Am J Anim Vet Sci* **11**:91–99 (2016). <https://doi.org/10.3844/ajavsp.2016.91.99>.
- 47 Whitley NC, Cazac D, Rude BJ, Jackson-O'Brien D and Parveen S, Use of a commercial probiotic supplement in meat goats. *J Anim Sci* **87**:723–728 (2009). <https://doi.org/10.2527/jas.2008-1031>.
- 48 Worku M, Adjei-Fremah S, Ekwemalor K, Asiamah E and Ismail H, 0130 growth and transcriptional profile analysis following oral probiotic supplementation in dairy cows. *J Anim Sci* **94**:61–61 (2016). <https://doi.org/10.2527/jam2016-0130>.
- 49 Deng Q, Odhiambo JF, Farooq U, Lam T, Dunn SM and Ametaj BN, Intravaginal probiotics modulated metabolic status and improved milk production and composition of transition dairy cows 1. *J Anim Sci* **94**:760–770 (2016). <https://doi.org/10.2527/jas.2015-9650>.
- 50 Ekwemalor K, Asiamah E, Osei B, Ismail H and Worku M, Evaluation of the effect of probiotic administration on gene expression in goat blood. *J Mol Biol Res* **7**:88 (2017). <https://doi.org/10.5539/jmbr.v7n1p88>.
- 51 Ding J, Zhou ZM, Ren LP and Meng QX, Effect of monensin and live yeast supplementation on growth performance, nutrient digestibility, carcass characteristics and ruminal fermentation parameters in lambs fed steam-flaked corn-based diets. *Asian-Australas J Anim Sci* **21**:547–554 (2008). <https://doi.org/10.5713/ajas.2008.70353>.
- 52 Puniya AK, Salem AZM, Kumar S, Dagar SS, Griffith GW, Puniya M *et al.*, Role of live microbial feed supplements with reference to anaerobic

- fungi in ruminant productivity: a review. *J Integr Agric* **14**:550–560 (2015). [https://doi.org/10.1016/S2095-3119\(14\)60837-6](https://doi.org/10.1016/S2095-3119(14)60837-6).
- 53 Elghandour MMY, Salem AZM, Castañeda JSM, Camacho LM, Kholif AE and Chagoyán JCV, Direct-fed microbes: a tool for improving the utilization of low quality roughages in ruminants. *J Integr Agric* **14**:526–533 (2015). [https://doi.org/10.1016/S2095-3119\(14\)60834-0](https://doi.org/10.1016/S2095-3119(14)60834-0).
  - 54 Nocek JE, Kautz WP, Leedle JAZ and Allman JG, Ruminant supplementation of direct-fed microbials on diurnal pH variation and in situ digestion in dairy cattle. *J Dairy Sci* **85**:429–433 (2002). [https://doi.org/10.3168/jds.S0022-0302\(02\)74091-5](https://doi.org/10.3168/jds.S0022-0302(02)74091-5).
  - 55 Kung L, Taylor CC, Lynch MP and Neylon JM, The effect of treating alfalfa with *Lactobacillus buchneri* 40788 on silage fermentation, aerobic stability, and nutritive value for lactating dairy cows. *J Dairy Sci* **86**:336–343 (2003). [https://doi.org/10.3168/jds.S0022-0302\(03\)73611-X](https://doi.org/10.3168/jds.S0022-0302(03)73611-X).
  - 56 Robinson AKP, Yeast products for growing and lactating dairy cattle: impact on rumen fermentation and performance. *Dairy Rev* **9**:1–4 (2002). <https://doi.org/10.3923/ajava.2012.366.375>.
  - 57 Stein DR, Allen DT, Perry EB, Bruner JC, Gates KW, Rehberger TG *et al.*, Effects of feeding propionibacteria to dairy cows on milk yield, milk components, and reproduction. *J Dairy Sci* **89**:111–125 (2006). [https://doi.org/10.3168/jds.S0022-0302\(06\)72074-4](https://doi.org/10.3168/jds.S0022-0302(06)72074-4).
  - 58 Weiss WP, Wyatt DJ and McKelvey TR, Effect of feeding propionibacteria on milk production by early lactation dairy cows. *J Dairy Sci* **91**:646–652 (2008). <https://doi.org/10.3168/jds.2007-0693>.
  - 59 Huntington GB, High-starch rations for ruminant production discussed. *Feedstuffs* **12**:13–23 (2000).
  - 60 van Soest PJ, *Nutritional Ecology of the Ruminant*. Cornell University Press, Ithaca (1994).
  - 61 Ghorbani GR, Morgavi DP, Beauchemin KA and Leedle JAZ, Effects of bacterial direct-fed microbials on ruminal fermentation, blood variables, and the microbial populations of feedlot cattle<sup>1,2</sup>. *J Anim Sci* **80**:1977–1985 (2002a). <https://doi.org/10.2527/2002.8071977x>.
  - 62 Chiquette J, Allison MJ and Rasmussen M, Use of *Prevotella bryantii* 25A and a commercial probiotic during subacute acidosis challenge in midlactation dairy cows. *J Dairy Sci* **95**:5985–5995 (2012). <https://doi.org/10.3168/jds.2012-5511>.
  - 63 Ghorbani GR, Morgavi DP, Beauchemin KA and Leedle JAZ, Effects of bacterial direct-fed microbials and yeast on site and extent of digestion, blood chemistry and subclinical ruminal acidosis in feedlot cattle. *J Anim Sci* **81**:1628–1640 (2003). <https://doi.org/10.2527/2003.8161628x>.
  - 64 Le OT, Dart PJ, Harper K, Zhang D, Schofield B, Callaghan MJ *et al.*, Effect of probiotic *Bacillus amyloliquefaciens* strain H57 on productivity and the incidence of diarrhoea in dairy calves. *Anim Prod Sci* **57**:912 (2017). <https://doi.org/10.1071/AN15776>.
  - 65 Nagaraja TG and Titgemeyer EC, Ruminal acidosis in beef cattle: the current microbiological and nutritional outlook. *J Dairy Sci* **90**:E17–E38 (2007). <https://doi.org/10.3168/jds.2006-478>.
  - 66 Rolfe RD, The role of probiotic cultures in the control of gastrointestinal health. *J Nutr* **130**:396S–402S (2000). <https://doi.org/10.1093/jn/130.2.396s>.
  - 67 La Ragione RM, Narbad A, Gasson MJ and Woodward MJ, In vivo characterization of *Lactobacillus johnsonii* F19785 for use as a defined competitive exclusion agent against bacterial pathogens in poultry. *Lett Appl Microbiol* **38**:197–205 (2004). <https://doi.org/10.1111/j.1472-765X.2004.01474.x>.
  - 68 Frizzo LS, Soto LP, Zbrun MV, Bertozzi E, Sequeira G, Armesto RR *et al.*, Lactic acid bacteria to improve growth performance in young calves fed milk replacer and spray-dried whey powder. *Anim Feed Sci Technol* **157**:159–167 (2010). <https://doi.org/10.1016/j.anifeeds.2010.03.005>.
  - 69 Forestier C, de Champs C, Vatoux C and Joly B, Probiotic activities of *Lactobacillus casei rhamnosus*: in vitro adherence to intestinal cells and antimicrobial properties. *Res Microbiol* **152**:167–173 (2001). [https://doi.org/10.1016/S0923-2508\(01\)01188-3](https://doi.org/10.1016/S0923-2508(01)01188-3).
  - 70 Fuller R, The chicken gut microflora and probiotic supplements. *J Poultry Sci* **38**:189–196 (2001). <https://doi.org/10.2141/jpsa.38.189>.
  - 71 Dicks LMT and Botes M, Probiotic lactic acid bacteria in the gastrointestinal tract: health benefits, safety and mode of action. *Benef Microbes* **1**:11–29 (2010). <https://doi.org/10.3920/BM2009.0012>.
  - 72 Cheikhoussef A, Pogori N, Chen W and Zhang H, Antimicrobial proteinaceous compounds obtained from bifidobacteria: from production to their application. *Int J Food Microbiol* **125**:215–222 (2008). <https://doi.org/10.1016/j.ijfoodmicro.2008.03.012>.
  - 73 Flynn S, van Sinderen D, Thornton GM, Holo H, Nes IF and Collins JK, Characterization of the genetic locus responsible for the production of ABP-118, a novel bacteriocin produced by the probiotic bacterium *Lactobacillus salivarius* subsp. *salivarius* UCC118. *Microbiology* **148**:973–984 (2002). <https://doi.org/10.1099/00221287-148-4-973>.
  - 74 Cotter PD, Hill C and Ross RP, Bacteriocins: developing innate immunity for food. *Nat Rev Microbiol* **3**:777–788 (2005). <https://doi.org/10.1038/nrmicro1273>.
  - 75 Rea MC, Clayton E, O'Connor PM, Shanahan F, Kiely B, Ross RP *et al.*, Antimicrobial activity of lactacin 3147 against clinical *Clostridium difficile* strains. *J Med Microbiol* **56**:940–946 (2007). <https://doi.org/10.1099/jmm.0.47085-0>.
  - 76 Matsuguchi T, Takagi A, Matsuzaki T, Nagaoka M, Ishikawa K, Yokokura T *et al.*, Lipoteichoic acids from *Lactobacillus* strains elicit strong tumor necrosis factor alpha-inducing activities in macrophages through toll-like receptor 2. *Clin Diagn Lab Immunol* **10**:259–266 (2003). <https://doi.org/10.1128/CDLI.10.2.259-266.2003>.
  - 77 Johnson-Henry KC, Hagen KE, Gordonpour M, Tompkins TA and Sherman PM, Surface-layer protein extracts from *Lactobacillus helveticus* inhibit enterohaemorrhagic *Escherichia coli* O157:H7 adhesion to epithelial cells. *Cell Microbiol* **9**:356–367 (2007). <https://doi.org/10.1111/j.1462-5822.2006.00791.x>.
  - 78 Bernet MF, Brassart D, Neeser JR and Servin AL, *Lactobacillus acidophilus* LA 1 binds to cultured human intestinal cell lines and inhibits cell attachment and cell invasion by enterovirulent bacteria. *Gut* **35**:483–489 (1994). <https://doi.org/10.1136/gut.35.4.483>.
  - 79 Waters CM and Bassler BL, Quorum sensing: cell-to-cell communication in bacteria. *Annu Rev Cell Dev Biol* **21**:319–346 (2005). <https://doi.org/10.1146/annurev.cellbio.21.012704.131001>.
  - 80 Hughes DT and Sperandio V, Inter-kingdom signalling: communication between bacteria and their hosts. *Nat Rev Microbiol* **6**:111–120 (2008). <https://doi.org/10.1038/nrmicro1836>.
  - 81 Medellín-Peña MJ, Wang H, Johnson R, Anand S and Griffiths MW, Probiotics affect virulence-related gene expression in *Escherichia coli* O157:H7. *Appl Environ Microbiol* **73**:4259–4267 (2007). <https://doi.org/10.1128/AEM.00159-07>.
  - 82 Lukic J, Jancic I, Mirkovic N, Bufan B, Djokic J, Milenkovic M *et al.*, *Lactococcus lactis* and *Lactobacillus salivarius* differently modulate early immunological response of Wistar rats co-administered with *Listeria monocytogenes*. *Benef Microbes* **8**:809–822 (2017). <https://doi.org/10.3920/BM2017.0007>.
  - 83 Dong Q, Zhang W, Guo L, Niu H, Liu Q and Wang X, Influence of *Lactobacillus plantarum* individually and in combination with low O2-MAP on the pathogenic potential of *Listeria monocytogenes* in cabbage. *Food Control* **107**:106765 (2020). <https://doi.org/10.1016/j.foodcont.2019.106765>.
  - 84 Bakr HA, Hassan MS, Giadinis ND, Panousis N, Ostojic-Andric D, El-Tawab A *et al.*, Effect of *Saccharomyces cerevisiae* supplementation on health and performance of dairy cows during transition and early lactation period. *Biotechnol Anim Husbandry Biotechnologija u Stoc* **31**:349–364 (2015). <https://doi.org/10.2298/bah1503349b>.
  - 85 Newbold CJ, Wallace RJ and McIntosh FM, Mode of action of the yeast *Saccharomyces cerevisiae* as a feed additive for ruminants. *Br J Nutr* **76**:249–261 (1996). <https://doi.org/10.1079/bjn19960029>.
  - 86 Bombá A, Nemcová R, Gancarciková S, Herich R, Guba P and Mudronová D, Improvement of the probiotic effect of microorganisms by their combination with maltodextrins, fructooligosaccharides and polyunsaturated fatty acids. *Br J Nutr* **88**:S95–S99 (2002). <https://doi.org/10.1079/bjn2002634>.
  - 87 Chaucheyras-Durand F, Walker ND and Bach A, Effects of active dry yeasts on the rumen microbial ecosystem: past, present and future. *Anim Feed Sci Technol* **145**:5–26 (2008). <https://doi.org/10.1016/j.anifeeds.2007.04.019>.
  - 88 Jouany JP, Optimizing rumen functions in the close-up transition period and early lactation to drive dry matter intake and energy balance in cows. *Anim Reprod Sci* **96**:250–264 (2006). <https://doi.org/10.1016/j.anireprosci.2006.08.005>.
  - 89 Aslan V, Thamsborg S, Jørgensen R and Basse A, Induced acute ruminal acidosis in goats treated with yeast (*Saccharomyces cerevisiae*) and bicarbonate. *Acta Vet Scand* **36**:65–77 (1995).
  - 90 Newman KE and Jacques KA, Microbial feed additives for pre-ruminants, in *Biotechnology in Animal Feeds and Animal Feeding*. Wiley-

- VCH Verlag GmbH, Weinheim, Germany, pp. 247–258 (2007). <https://doi.org/10.1002/9783527615353.ch12>.
- 91 Niemiłowski M, Schollenberger A and Klucinski W, Mucosal immunity and the bovine entero-mammary link: evolutionary established dialogue between antigen and arms of immune system, in *Biology of Growing Animals*. Elsevier, **2**:293–313 (2005). [https://doi.org/10.1016/S1877-1823\(09\)70046-3](https://doi.org/10.1016/S1877-1823(09)70046-3).
  - 92 Thormar H, Patented non-antibiotic agents as animal feed additives. *Recent Pat Food Nutr Agric* **4**:155–168 (2012). <https://doi.org/10.2174/2212798411204020155>.
  - 93 Donovan DC, Franklin ST, Chase CCL and Hippen AR, Growth and health of Holstein calves fed milk replacers supplemented with antibiotics or enteroguard. *J Dairy Sci* **85**:947–950 (2002). [https://doi.org/10.3168/jds.S0022-0302\(02\)74153-2](https://doi.org/10.3168/jds.S0022-0302(02)74153-2).
  - 94 Isolauri E, Sütas Y, Kankaanpää P, Arvilommi H and Salminen S, Probiotics: effects on immunity. *Am J Clin Nutr* **73**:444s–450s (2001). <https://doi.org/10.1093/ajcn/73.2.444s>.
  - 95 Murphy K, Paul T, Mark W and Charles J, *Janeway's Immunobiology*, 8th edn. Garland Science, New York, NY (2012).
  - 96 Adjei-Fremah S, Ekwemalor K, Asiamah EK, Ismail H, Ibrahim S and Worku M, Effect of probiotic supplementation on growth and global gene expression in dairy cows. *J Appl Anim Res* **46**:257–263 (2018). <https://doi.org/10.1080/09712119.2017.1292913>.
  - 97 Guedes CM, Gonçalves D, Rodrigues MAM and Dias-da-Silva A, Effects of a *Saccharomyces cerevisiae* yeast on ruminal fermentation and fibre degradation of maize silages in cows. *Anim Feed Sci Technol* **145**:27–40 (2008). <https://doi.org/10.1016/j.anifeedsci.2007.06.037>.
  - 98 Hasunuma T, Kawashima K, Nakayama H, Murakami T, Kanagawa H, Ishii T et al., Effect of cellooligosaccharide or synbiotic feeding on growth performance, fecal condition and hormone concentrations in Holstein calves. *Anim Sci J* **82**:543–548 (2011). <https://doi.org/10.1111/j.1740-0929.2010.00861.x>.
  - 99 O'Hara AM and Shanahan F, Gut microbiota: Mining for therapeutic potential. *Clin Gastroenterol Hepatol* **5**:274–284 (2007). <https://doi.org/10.1016/j.cgh.2006.12.009>.
  - 100 Poppy GD, Rabiee AR, Lean IJ, Sanchez WK, Dorton KL and Morley PS, A meta-analysis of the effects of feeding yeast culture produced by anaerobic fermentation of *Saccharomyces cerevisiae* on milk production of lactating dairy cows. *J Dairy Sci* **95**:6027–6041 (2012). <https://doi.org/10.3168/jds.2012-5577>.
  - 101 Saleem AM, Zounouny AI and Singer AM, Growth performance, nutrients digestibility, and blood metabolites of lambs fed diets supplemented with probiotics during pre- and post-weaning period. *Asian-Australas J Anim Sci* **30**:523–530 (2017). <https://doi.org/10.5713/ajas.16.0691>.
  - 102 Lesmeister KE, Heinrichs AJ and Gabler MT, Effects of supplemental yeast (*Saccharomyces cerevisiae*) culture on rumen development, growth characteristics, and blood parameters in neonatal dairy calves. *J Dairy Sci* **87**:1832–1839 (2004). [https://doi.org/10.3168/jds.S0022-0302\(04\)73340-8](https://doi.org/10.3168/jds.S0022-0302(04)73340-8).
  - 103 Hilla H, El-Sayaad G and Abdella M, Effect of growth promoters (probiotics) supplementation on performance, rumen activity and some blood constituents in growing lambs. *Arch Anim Breed* **54**:607–617 (2011). <https://doi.org/10.5194/aab-54-607-2011>.
  - 104 Mudgal V and Baghel R, Effect of probiotic supplementation on growth performance of pre-ruminant buffalo (*Bubalus bubalis*) calves. *Buffalo Bull* **29**:37–39 (2010).
  - 105 Reuben RC, Roy PC, Sarkar SL, Alam RU and Jahid IK, Isolation, characterization, and assessment of lactic acid bacteria toward their selection as poultry probiotics. *BMC Microbiol* **19**:253 (2019). <https://doi.org/10.1186/s12866-019-1626-0>.
  - 106 Reuben RC, Roy PC, Sarkar SL, Rubayet UI Alam ASM and Jahid IK, Characterization and evaluation of lactic acid bacteria from indigenous raw milk for potential probiotic properties. *J Dairy Sci* **103**:1223–1237 (2020). <https://doi.org/10.3168/jds.2019-17092>.
  - 107 Song D, Ibrahim S and Hayek S, Recent application of probiotics in food and agricultural science, in *Probiotics*. InTech, London, UK (2012). <https://doi.org/10.5772/50121>.
  - 108 Sretenovic L, Petrovic M, Aleksic S, Pantelic V, Katic V, Bogdanovic V et al., Influence of yeast, probiotics and enzymes in rations on dairy cows performances during transition. *Biotechnol Anim Husband* **24**:33–43 (2008). <https://doi.org/10.2298/bah0806033s>.
  - 109 Xu H, Huang W, Hou Q, Kwok L Y, Sun Z, Ma H et al., The effects of probiotics administration on the milk production, milk components and fecal bacteria microbiota of dairy cows. *Sci Bull* **62**:767–774 (2017). <https://doi.org/10.1016/j.scib.2017.04.019>.
  - 110 Yu P, Huber JT, Theurer CB, Chen KH, Nussio LG and Wu Z, Effect of steam-flaked or steam-rolled corn with or without *Aspergillus oryzae* in the diet on performance of dairy cows fed during hot weather. *J Dairy Sci* **80**:3293–3297 (1997). [https://doi.org/10.3168/jds.S0022-0302\(97\)76304-5](https://doi.org/10.3168/jds.S0022-0302(97)76304-5).
  - 111 Qiao GH, Shan AS, Ma N, Ma QQ and Sun ZW, Effect of supplemental *Bacillus* cultures on rumen fermentation and milk yield in Chinese Holstein cows. *J Anim Physiol Anim Nutr (Berl)* **94**:429–436 (2010). <https://doi.org/10.1111/j.1439-0396.2009.00926.x>.
  - 112 Sun P, Wang JQ and Zhang HT, Effects of supplementation of *Bacillus subtilis* natto Na and N1 strains on rumen development in dairy calves. *Anim Feed Sci Technol* **164**:154–160 (2011). <https://doi.org/10.1016/j.anifeedsci.2011.01.003>.
  - 113 Stella AV, Paratte R, Valnegri L, Cigalino G, Soncini G, Chevaux E et al., Effect of administration of live *Saccharomyces cerevisiae* on milk production, milk composition, blood metabolites, and faecal flora in early lactating dairy goats. *Small Rumin Res* **67**:7–13 (2007). <https://doi.org/10.1016/j.smallrumres.2005.08.024>.
  - 114 Desnoyers M, Giger-Reverdin S, Bertin G, Duvaux-Ponter C and Sauvant D, Meta-analysis of the influence of *Saccharomyces cerevisiae* supplementation on ruminal parameters and milk production of ruminants. *J Dairy Sci* **92**:1620–1632 (2009). <https://doi.org/10.3168/jds.2008-1414>.
  - 115 Maamourai O, Selmi H and M'Hamdi N, Effects of yeast (*Saccharomyces cerevisiae*) feed supplement on milk production and its composition in Tunisian Holstein Friesian cows. *Sci Agric Bohem* **45**:170–174 (2014). <https://doi.org/10.2478/sab-2014-0104>.
  - 116 Kritas SK, Govaris A, Christodoulopoulos G and Burriel AR, Effect of *Bacillus licheniformis* and *Bacillus subtilis* supplementation of ewe's feed on sheep milk production and young lamb mortality. *J Vet Med Ser A Physiol Pathol Clin Med* **53**:170–173 (2006). <https://doi.org/10.1111/j.1439-0442.2006.00815.x>.
  - 117 Arowolo MA and He J, Use of probiotics and botanical extracts to improve ruminant production in the tropics: a review. *Anim Nutr* **4**:241–249 (2018). <https://doi.org/10.1016/j.aninu.2018.04.010>.
  - 118 El-Waziry AM and Ibrahim HR, Effect of *Saccharomyces cerevisiae* of yeast on fiber digestion in sheep fed Berseem (*Trifolium alexandrinum*) Hay and cellulase activity. *Aust J Basic Appl Sci* **1**:379–385 (2007).
  - 119 Kamel HEM, Sekine J, El-Waziry AM and Yacout MHM, Effect of *Saccharomyces cerevisiae* on the synchronization of organic matter and nitrogen degradation kinetics and microbial nitrogen synthesis in sheep fed Berseem hay (*Trifolium alexandrinum*). *Small Rumin Res* **52**:211–216 (2004). <https://doi.org/10.1016/j.smallrumres.2003.06.001>.
  - 120 Callaway ES and Martin SA, Effects of a *Saccharomyces cerevisiae* culture on ruminal bacteria that utilize lactate and digest cellulose. *J Dairy Sci* **80**:2035–2044 (1997). [https://doi.org/10.3168/jds.S0022-0302\(97\)76148-4](https://doi.org/10.3168/jds.S0022-0302(97)76148-4).
  - 121 Tungland BC and Meyer D, Nondigestible oligo- and polysaccharides (dietary fiber): their physiology and role in human health and food. *Compr Rev Food Sci Food Saf* **1**:90–109 (2002). <https://doi.org/10.1111/j.1541-4337.2002.tb00009.x>.
  - 122 Moore KJ and Hatfield RD, Carbohydrates and forage quality, in *Forage Quality, Evaluation, and Utilization*. American Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Madison, WI, pp. 229–280 (2015). <https://doi.org/10.2134/1994.foragequality.c6>.
  - 123 Arcos-García JL, Castrejón FA, Mendoza GD and Pérez-Gavilán EP, Effect of two commercial yeast cultures with *Saccharomyces cerevisiae* on ruminal fermentation and digestion in sheep fed sugar cane tops. *Livest Prod Sci* **63**:153–157 (2000). [https://doi.org/10.1016/S0301-6226\(99\)00116-5](https://doi.org/10.1016/S0301-6226(99)00116-5).
  - 124 Haddad SG and Goussou SN, Effect of yeast culture supplementation on nutrient intake, digestibility and growth performance of Awassi lambs. *Anim Feed Sci Technol* **118**:343–348 (2005). <https://doi.org/10.1016/j.anifeedsci.2004.10.003>.
  - 125 Boyd J, West JW and Bernard JK, Effects of the addition of direct-fed microbials and glycerol to the diet of lactating dairy cows on milk yield and apparent efficiency of yield. *J Dairy Sci* **94**:4616–4622 (2011).
  - 126 Azzaz HE-DH, Farahat ESA and Murad HA, Impact of microbial feed supplements on the productive performance of lactating nubian goats. *Global Vet* **14**:241–249 (2015).

- 127 Mousa K, El-Malky O, Komonna O and Rashwan SE, Effect of live dried yeast supplementation on digestion coefficients, some rumen fermentation, blood constituents and some reproductive and productive parameters in Rahmani sheep. *J Anim Sci* **8**:291–303 (2012). <https://doi.org/10.1007/s11250-012-0257-3>.
- 128 Stover MG, Watson RR and Collier RJ, Pre-and probiotic supplementation in ruminant livestock production, in *Probiotics, Prebiotics, and Synbiotics: Bioactive Foods in Health Promotion*. Elsevier Inc., London, UK, pp. 25–36 (2016). <https://doi.org/10.1016/B978-0-12-802189-7.00002-2>.
- 129 Sargeant JM, Amezcua MR, Rajic A and Waddell L, Pre-harvest interventions to reduce the shedding of *E. coli* O157 in the faeces of weaned domestic ruminants: a systematic review. *Zoonoses Public Health* **54**:260–277 (2007). <https://doi.org/10.1111/j.1863-2378.2007.01059.x>.
- 130 Wisener LV, Sargeant JM, O'Connor AM, Faires MC and Glass-Kaasta SK, The use of direct-fed microbials to reduce shedding of *Escherichia coli* O157 in beef cattle: a systematic review and meta-analysis. *Zoonoses Public Health* **62**:75–89 (2015). <https://doi.org/10.1111/zph.12112>.
- 131 Younts-Dahl SM, Osborn GD, Galyean ML, Rivera JD, Loneragan GH and Brashears MM, Reduction of *Escherichia coli* O157 in finishing beef cattle by various doses of *Lactobacillus acidophilus* in direct-fed microbials. *J Food Prot* **68**:6–10 (2005). <https://doi.org/10.4315/0362-028X-68.1.6>.
- 132 Jatkauskas J and Vrotniakiene V, Effects of probiotic dietary supplementation on diarrhoea patterns, faecal microbiota and performance of early weaned calves. *Vet Med (Praha)* **55**:494–503 (2010). <https://doi.org/10.17221/2939-VETMED>.
- 133 Galip N, Effect of supplemental yeast culture and sodium bicarbonate on ruminal fermentation and blood variables in rams. *J Anim Physiol Anim Nutr (Berl)* **90**:446–452 (2006). <https://doi.org/10.1111/j.1439-0396.2006.00625.x>.
- 134 Antunović Z, Šperanda M, Liker B, Šerić V, Senčić D, Domaćinović M *et al.*, Influence of feeding the probiotic Pioneer PDFM® to growing lambs on performances and blood composition. *Acta Vet Brno* **55**:287–300 (2005). <https://doi.org/10.2298/AVB0504287A>.
- 135 Bruno RGS, Rutigliano HM, Cerri RL, Robinson PH and Santos JEP, Effect of feeding saccharomyces *Cerevisiae* on performance of dairy cows during summer heat stress. *Anim Feed Sci Technol* **150**:175–186 (2009). <https://doi.org/10.1016/j.anifeeds.2008.09.001>.
- 136 Sayed A, Studies on the influences of pronifer as a probiotic on the clinical, hematological and biochemical status of goat kids. *Assiut Vet Med J* **99**:131–143 (2003).
- 137 Samanya M and Yamauchi K, Histological alterations of intestinal villi in chickens fed dried *Bacillus subtilis* var. natto. *Comp Biochem Physiol - A Mol Integr Physiol* **133**:95–104 (2002). [https://doi.org/10.1016/S1095-6433\(02\)00121-6](https://doi.org/10.1016/S1095-6433(02)00121-6).
- 138 Cetin N, Guclu BK and Cetin E, The effects of probiotic and mannanoligosaccharide on some haematological and immunological parameters in turkeys. *J Vet Med A Physiol Pathol Clin Med* **52**:263–267 (2005). <https://doi.org/10.1111/j.1439-0442.2005.00736.x>.
- 139 Deraz S, Synergetic effects of multispecies probiotic supplementation on certain blood parameters and serum biochemical profile of broiler chickens. *J Anim Health Prod* **2003**:27–34 (2018). <https://doi.org/10.1017/s1752756200013442>.
- 140 LaFleur B and LaFleur D, *Exploring Medical Language: A Student-Directed Approach*, 10th edn. Myrna LaFleur Brooks, Phoenix, USA (2017).
- 141 Reuben RC, Sarkar SL, Ibanat H, Ali AH, Roy RC and Iqbal IJ, Novel multi-strain probiotics reduces *Pasteurella multocida* induced fowl cholera mortality in broilers. *Sci Rep* **11**:8885 (2021). <https://doi.org/10.1038/s41598-021-88299-0>.
- 142 Abas I, Kutay HC, Kahraman R, Tokar NY, Özçelik D, Ates F *et al.*, Effects of organic acid and bacterial direct-feed microbial on fattening performance of Kivircik-male yearling lambs. *Pak J Nutr* **6**:149–154 (2007). <https://doi.org/10.3923/pjn.2007.149.154>.
- 143 Chiofalo V, Liotta L and Chiofalo B, Effects of the administration of *Lactobacilli* on body growth and on the metabolic profile in growing Maltese goat kids. *Reprod Nutr Dev* **44**:449–457 (2004). <https://doi.org/10.1051/rnd:2004051>.
- 144 Panda AK, Rama Rao SV, Raju MVLN and Sharma SR, Dietary supplementation of *Lactobacillus sporogenes* on performance and serum Biochemico - lipid profile of broiler chickens. *J Poultry Sci* **43**:235–240 (2006). <https://doi.org/10.2141/jpsa.43.235>.
- 145 Fukushima M and Nakano M, The effect of a probiotic on faecal and liver lipid classes in rats. *Br J Nutr* **73**:701–710 (1995). <https://doi.org/10.1079/bjn19950074>.
- 146 Mohan B, Kadirvel R, Natarajan A and Bhaskaran M, Effect of probiotic supplementation on growth, nitrogen utilisation and serum cholesterol in broilers. *Br Poultry Sci* **37**:395–401 (1996). <https://doi.org/10.1080/00071669608417870>.
- 147 Alkhalif A, Alhaj M and Al-Homidan I, Influence of probiotic supplementation on blood parameters and growth performance in broiler chickens. *Saudi J Biol Sci* **17**:219–225 (2010). <https://doi.org/10.1016/j.sjbs.2010.04.005>.
- 148 Surono IS, In vitro probiotic properties of indigenous dadih lactic acid bacteria. *Asian-Australas J Anim Sci* **16**:726–731 (2003). <https://doi.org/10.5713/ajas.2003.726>.
- 149 Pulina G, Francesconi AHD, Stefanon B, Sevi A, Calamari L, Lacetera N *et al.*, Sustainable ruminant production to help feed the planet. *Ital J Anim Sci* **16**:140–171 (2017). <https://doi.org/10.1080/1828051X.2016.1260500>.
- 150 Lambin EF, Global land availability: Malthus versus Riccardo. *Global Food Secur* **1**:83–87 (2012).
- 151 Goldberg AM, Farm animal welfare and human health. *Curr Environ Health Rep* **3**:313–321 (2016).
- 152 Atkinson C, Organic animal husbandry in Europe – current and future challenges, in *Organic Animal Husbandry across the World: Towards an Action Plan for Development of Organic Animal Husbandry*, ed. by Schmid O, Chander M, Escosteguy A and Frueh B. IAHAIFOAM Animal Husbandry Alliance, Schweiz, pp. 25–27 (2014).
- 153 Kowalski ZM, Górka P, Schlagheck A, Jagusiak W, Micek P and Strzetelski J, Performance of Holstein calves fed milk-replacer and starter mixture supplemented with probiotic feed additive. *J Anim Feed Sci* **18**:399–411 (2009). <https://doi.org/10.22358/jafs/66409/2009>.
- 154 Salazar L, Nero L, Campos-Galvão M, Cortinhas C, Acedo T, Tamassia L *et al.*, Effect of oselected feed additives to improve growth and health of dairy calves. *PLoS One* **14**:e0216066 (2019). <https://doi.org/10.1371/journal.pone.0216066>.
- 155 Roodposhti PM and Dabiri N, Effects of probiotic and prebiotic on average daily gain, fecal shedding of *Escherichia coli*, and immune system status in newborn female calves. *Asian-Australas J Anim Sci* **25**:1255–1261 (2012). <https://doi.org/10.5713/ajas.2011.11312>.
- 156 Kafizadeh F, Payandeh S, Gómez-Cortés P, Ghadimi D, Schiavone A and Martínez Marín AL, Effects of probiotic supplementation on milk production, blood metabolite profile and enzyme activities of ewes during lactation. *Ital J Anim Sci* **18**:134–139 (2019). <https://doi.org/10.1080/1828051X.2018.1496040>.
- 157 Yasuda K, Hashikawa S, Sakamoto H, Tomita Y, Shibata S and Fukata T, A new synbiotic consisting of *Lactobacillus casei* subsp. *casei* and dextran improves milk production in holstein dairy cows. *J Vet Med Sci* **69**:205–208 (2007). <https://doi.org/10.1292/jvms.69.205>.
- 158 Stęczyński K and Kokoszyński D, Effect of probiotic preparations (EM) and sex on morphometric characteristics of the digestive system and leg bones, and caecal microflora in broiler chickens. *J Appl Anim Res* **48**:45–50 (2020). <https://doi.org/10.1080/09712119.2020.1718680>.
- 159 Tristant D and Moran CA, The efficacy of feeding a live probiotic yeast, Yea-Sacc®, on the performance of lactating dairy cows. *J Appl Anim Nutr* **3**:1–16 (2015). <https://doi.org/10.1017/jan.2015.10>.
- 160 Beharka AA, Nagaraja TG and Morrill JL, Performance and ruminal function development of young calves fed diets with *Aspergillus oryzae* fermentation extract. *J Dairy Sci* **74**:4326–4336 (1991). [https://doi.org/10.3168/jds.S0022-0302\(91\)78628-1](https://doi.org/10.3168/jds.S0022-0302(91)78628-1).
- 161 Cruywagen CW, Jordan I and Venter L, Effect of *Lactobacillus acidophilus* supplementation of milk replacer on preweaning performance of calves. *J Dairy Sci* **79**:483–486 (1996).
- 162 Abe F, Ishibashi N and Shimamura S, Effect of administration of bifidobacteria and lactic acid bacteria to newborn calves and piglets. *J Dairy Sci* **78**:2838–2846 (1995). [https://doi.org/10.3168/jds.S0022-0302\(95\)76914-4](https://doi.org/10.3168/jds.S0022-0302(95)76914-4).
- 163 Malik R and Bandla S, Effect of source and dose of probiotics and exogenous fibrolytic enzymes (EFE) on intake, feed efficiency, and growth of male buffalo (*Bubalus bubalis*) calves. *Trop Anim Health Prod* **42**:1263–1269 (2010). <https://doi.org/10.1007/s11250-010-9559-5>.
- 164 Nocek JE and Kautz WP, Direct-fed microbial supplementation on ruminal digestion, health, and performance of pre- and postpartum

- dairy cattle. *J Dairy Sci* **89**:260–266 (2006). [https://doi.org/10.3168/jds.S0022-0302\(06\)72090-2](https://doi.org/10.3168/jds.S0022-0302(06)72090-2).
- 165 Raeth-Knight ML, Linn JG and Jung HG, Effect of direct-fed microbials on performance, diet digestibility, and rumen characteristics of Holstein dairy cows. *J Dairy Sci* **90**:1802–1809 (2007). <https://doi.org/10.3168/jds.2006-643>.
  - 166 Keyser SA, McMeniman JP, Smith DR, MacDonald JC and Galyean ML, Effects of *Saccharomyces cerevisiae* subspecies *boulardii* CNCM I-1079 on feed intake by healthy beef cattle treated with florfenicol and on health and performance of newly received beef heifers. *J Anim Sci* **85**:1264–1273 (2007). <https://doi.org/10.2527/jas.2006-751>.
  - 167 Lehloenyha KV, Krehblel CR, Mertz KJ, Rehberger TG and Spicer LJ, Effects of propionibacteria and yeast culture fed to steers on nutrient intake and site and extent of digestion. *J Dairy Sci* **91**:653–662 (2008). <https://doi.org/10.3168/jds.2007-0474>.
  - 168 Lascano GJ, Heinrichs AJ and Tricarico JM, Substitution of starch by soluble fiber and *Saccharomyces cerevisiae* dose response on nutrient digestion and blood metabolites for precision-fed dairy heifers 1. *J Dairy Sci* **95**:3298–3309 (2012). <https://doi.org/10.3168/jds.2011-5047>.
  - 169 Arakaki LC, Stahringer RC, Garrett JE and Dehority BA, The effects of feeding monensin and yeast culture, alone or in combination, on the concentration and generic composition of rumen protozoa in steers fed on low-quality pasture supplemented with increasing levels of concentrate. *Anim Feed Sci Technol* **84**:121–127 (2000). [https://doi.org/10.1016/S0377-8401\(00\)00108-5](https://doi.org/10.1016/S0377-8401(00)00108-5).
  - 170 Miller-Webster T, Hoover WH, Holt M and Nocek JE, Influence of yeast culture on ruminal microbial metabolism in continuous culture. *J Dairy Sci* **85**:2009–2014 (2002). [https://doi.org/10.3168/jds.S0022-0302\(02\)74277-X](https://doi.org/10.3168/jds.S0022-0302(02)74277-X).
  - 171 Pinloche E, McEwan N, Marden J-P, Bayourthe C, Auclair E and Newbold CJ, The effects of a probiotic yeast on the bacterial diversity and population structure in the rumen of cattle. *PLoS One* **8**:e67824 (2013). <https://doi.org/10.1371/journal.pone.0067824>.
  - 172 Uyen Y, Shigemori S and Shimosato T, Effect of probiotics/prebiotics on cattle health and productivity. *Microbes Environ. Japanese Society of Microbial Ecology* **30**:126–132 (2015). <https://doi.org/10.1264/jsm2.ME14176>.
  - 173 Lee SS, Ha JK and Cheng KJ, Influence of an anaerobic fungal culture administration on in vivo ruminal fermentation and nutrient digestion. *Anim Feed Sci Technol* **88**:201–217 (2000). [https://doi.org/10.1016/S0377-8401\(00\)00216-9](https://doi.org/10.1016/S0377-8401(00)00216-9).
  - 174 Saxena S, Sehgal JP, Puniya AK and Singh K, Effect of administration of rumen fungi on production performance of lactating buffaloes. *Benef Microbes* **1**:183–188 (2010). <https://doi.org/10.3920/BM2009.0018>.
  - 175 Tripathi VK, Sehgal JP, Puniya AK and Singh K, Effect of administration of anaerobic fungi isolated from cattle and wild blue bull (*Boselaphus tragocamelus*) on growth rate and fibre utilization in buffalo calves. *Arch Anim Nutr* **61**:416–423 (2007). <https://doi.org/10.1080/17450390701556759>.
  - 176 Merrill ML, Bohnert DW, Harmon DL, Craig AM and Schrick FN, The ability of a yeast-derived cell wall preparation to minimize the toxic effects of high-ergot alkaloid tall fescue straw in beef cattle. *J Anim Sci* **85**:2596–2605 (2007). <https://doi.org/10.2527/jas.2007-0075>.
  - 177 Lu Q, Wu J, Wang M, Zhou C, Han X, Odongo EN et al., Effects of dietary addition of cellulase and a *Saccharomyces cerevisiae* fermentation product on nutrient digestibility, rumen fermentation and enteric methane emissions in growing goats. *Arch Anim Nutr* **70**:224–238 (2016). <https://doi.org/10.1080/1745039X.2016.1163002>.
  - 178 Kumar DS, Prasad CS and Prasad RMV, Effect of yeast culture (*Saccharomyces cerevisiae*) on ruminal microbial population in buffalo bulls. *Buff Bull* **32**:209–215 (2013).
  - 179 Jatkauskas J and Vrotniakien V, Effect of *L. plantarum*, *pediococcus acidilactici*, *enterococcus faecium* and *L. lactis* microbial supplementation of grass silage on the fermentation characteristics in rumen of dairy cows | semantic scholar. *Vet Zootech* **40**:29–34 (2007).
  - 180 Timmerman HM, Mulder L, Everts H, Van Espen DC, van der Wal E, Klaassen G et al., Health and growth of veal calves fed milk replacers with or without probiotics. *J Dairy Sci* **88**:2154–2165 (2005). [https://doi.org/10.3168/jds.S0022-0302\(05\)72891-5](https://doi.org/10.3168/jds.S0022-0302(05)72891-5).
  - 181 Jukna C, Jukna V and Imkus S, The influence of probiotic Lactomycolin on the growth of calves. *Vet Zootech* **6**:65–69 (2004).
  - 182 Das KC and Singh AS, Economics of feeding lactic acid bacteria fed on grain-based and grainless calf starter in calves. *Indian J Anim Res* **37**:44–47 (2002).
  - 183 Isik M, Ekimler F, O'zen N and Firat M, Effects of using probiotics on the growth performance and health of dairy calves | request PDF. *Turkish J Vet Anim Sci* **28**:63–69 (2004).
  - 184 Bilik K, Strzetelski J, Osiegowski S and Milewski J, Effect of various probiotics on cow performance and milk composition. *Rocz Nauk Zootech* **27**:169–177 (2000).
  - 185 Lema M, Williams L and Rao DR, Reduction of fecal shedding of enterohemorrhagic *Escherichia coli* O157:H7 in lambs by feeding microbial feed supplement. *Small Rumin Res* **39**:31–39 (2001). [https://doi.org/10.1016/S0921-4488\(00\)00168-1](https://doi.org/10.1016/S0921-4488(00)00168-1).
  - 186 Elam NA, Gleghorn JF, Rivera JD, Galyean ML, Defoor PJ, Brashears MM et al., Effects of live cultures of *Lactobacillus acidophilus* (strains NP45 and NP51) and *Propionibacterium freudenreichii* on performance, carcass, and intestinal characteristics, and *Escherichia coli* strain O157 shedding of finishing beef steers. *J Anim Sci* **81**:2686–2698 (2003). <https://doi.org/10.2527/2003.81112686x>.
  - 187 Roos TB, Tabeleão VC, Dümmler LA, Schwegler E, Goulart MA, Moura SV et al., Effect of *Bacillus cereus* var. *Toyo* and *Saccharomyces boulardii* on the immune response of sheep to vaccines. *Food Agric Immunol* **21**:113–118 (2010). <https://doi.org/10.1080/09540100903443691>.
  - 188 Arthur TM, Bosilevac JM, Kalchayanand N, Wells JE, Shackelford SD, Wheeler TL et al., Evaluation of a direct-fed microbial product effect on the prevalence and load of *Escherichia coli* O157:H7 in feedlot cattle. *J Food Prot* **73**:366–371 (2010). <https://doi.org/10.4315/0362-028X-73.2.366>.
  - 189 Görgülü M, Siuta A, Yurtseven S, Öngel E and Kutlu HR, Effect of probiotics on growing performance and health of calves. *Cuba J Agric Sci* **37**:125–129 (2003). <https://doi.org/10.3923/pjbs.2003.651.654>.
  - 190 Badié A, Moosakhani F, Hamidi A and Sami M, The effect of Protexin on prevention of ileocecal infection by *Mycobacterium avium* subspecies *paratuberculosis* in dairy calves. *J Dairy Sci* **96**:6535–6538 (2013). <https://doi.org/10.3168/jds.2012-5535>.
  - 191 Younts-Dahl SM, Galyean ML, Loneragan GH, Elam NA and Brashears MM, Dietary supplementation with *Lactobacillus*- and *Propionibacterium*-based direct-fed microbials and prevalence of *Escherichia coli* O157 in beef feedlot cattle and on hides at harvest. *J Food Prot* **67**:889–893 (2004). <https://doi.org/10.4315/0362-028X-67.5.889>.
  - 192 Brashears MM, Galyean ML, Loneragan GH, Mann JE and Killinger-Mann K, Prevalence of *Escherichia coli* O157:H7 and performance by beef feedlot cattle given *Lactobacillus* direct-fed microbials. *J Food Prot* **66**:748–754 (2003). <https://doi.org/10.4315/0362-028X-66.5.748>.
  - 193 Ohya T, Akiba M and Ito H, Use of a trial probiotic product in calves experimentally infected with *Escherichia coli* O157. *Jpn Agric Res Q* **35**:189–194 (2001). <https://doi.org/10.6090/jarq.35.189>.
  - 194 Ohya T, Marubashi T and Ito H, Significance of fecal volatile fatty acids in shedding of *Escherichia coli* O157 from calves: experimental infection and preliminary use of a probiotic product. *J Vet Med Sci* **62**:1151–1155 (2000). <https://doi.org/10.1292/jvms.62.1151>.
  - 195 Abu-Tarboush HM, Al-Saiady MY and Keir El-Din AH, Evaluation of diet containing *Lactobacilli* on performance, fecal coliform, and *Lactobacilli* of young dairy calves. *Anim Feed Sci Technol* **57**:39–49 (1996). [https://doi.org/10.1016/0377-8401\(95\)00850-0](https://doi.org/10.1016/0377-8401(95)00850-0).
  - 196 Wehnes CA, Novak KN, Patskevich V, Shields DR, Coalson JA, Smith AH et al., Benefits of supplementation of an electrolyte scour treatment with a *Bacillus*-based direct-fed microbial for calves. *Probiotics Antimicrob Proteins* **1**:36–44 (2009). <https://doi.org/10.1007/s12602-008-9004-5>.