

Potential impact of prickly pear cactus flour and Salix babylonica extract on cecal fermentation and methane production in horses


**A. Parra-Garcia, A. Z. M. Salem,
M. M. Y. Elghandour, L. M. Camacho &
N. E. Odongo**

Agroforestry Systems

An International Journal incorporating
Agroforestry Forum

ISSN 0167-4366
Volume 92
Number 5

Agroforest Syst (2018) 92:1145-1154
DOI 10.1007/s10457-016-0051-8



Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media Dordrecht. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Potential impact of prickly pear cactus flour and *Salix babylonica* extract on cecal fermentation and methane production in horses

A. Parra-Garcia · A. Z. M. Salem ·
M. M. Y. Elghandour · L. M. Camacho ·
N. E. Odongo

Received: 15 April 2016 / Accepted: 15 November 2016 / Published online: 23 November 2016
© Springer Science+Business Media Dordrecht 2016

Abstract The cecal gas (GP) and methane (CH₄) production and cecal fermentation kinetics when corn grain (CG) was replaced with prickly cactus (PC) in a horse's diet at different levels of *Salix babylonica* (SB) extract was investigated. Three total mixed rations where CG was replaced with PC at three levels (/kg): 0 g (Control), 75 g (PC75) or 150 g (PC150) were prepared and SB extract added at four levels: 0, 0.6, 1.2 and 1.8 mL/g dry matter (DM) of substrates. No ration type × SB extract dose interaction was observed ($P > 0.05$) for GP kinetics and CH₄ production. Increasing the level of PC in the ration quadratically increased ($P < 0.01$) the asymptotic GP and decreased ($P < 0.01$) the rate and lag time of GP. Increasing the level of PC in the ration, increased GP values ($P < 0.05$). Increasing the level of SB extract linearly decreased ($P = 0.001$) the lag time of GP of

all diets without affecting the asymptotic GP or the rate of GP. Ration type and SB level had no effect ($P > 0.05$) on CH₄ production; however, at 36 h of incubation, SB extract decreased CH₄ production. The rations PC75 and PC150 increased cecal pH compared with the control ration. The PC150 ration had the highest ($P < 0.05$) DM degradability, short chain fatty acids production, and gas yield after 24 h of incubation, with no effect ($P > 0.05$) of SB inclusion on all investigated fermentation kinetic parameters. It is concluded that increasing the level of PC in the diet of horse and replacing CG up to 60%, increased GP and improved cecal fermentation kinetics without affecting CH₄ production. Inclusion of *S. babylonica* extract in the tested rations had weak effects on fermentation kinetics although it decreased the lag time of GP.

A. Parra-Garcia · A. Z. M. Salem (✉) ·
M. M. Y. Elghandour
Facultad de Medicina Veterinaria y Zootecnia,
Universidad Autónoma del Estado de México, Toluca,
Mexico
e-mail: asalem70@yahoo.com

L. M. Camacho
Unidad Académica de Medicina Veterinaria y Zootecnia,
Universidad Autónoma de Guerrero, Cd. Altamirano-
Iguala, Chilpancingo, Guerrero, Mexico

N. E. Odongo
Department of Animal Sciences, School of Agriculture,
Pwani University, P. O. Box 195-80108, Kilifi, Kenya

Keywords Cecal fermentation · Corn grain · Horse nutrition · Prickly pear cactus · *Salix babylonica* extract

Introduction

Agricultural byproducts produced worldwide during different agricultural practices are nutrients-rich feed ingredients with a huge potential to be used in ruminant nutrition (Ahmed et al. 2015; Elghandour et al. 2016a). However, in many developing countries,

agriculture byproducts are not adequately utilized cause environmental problems resulting in their burning in the field. Such materials can be used as a cleaner product of animal feed and environmental conservation (Elghandour et al. 2016b, c). The livestock sector is suffering from feed shortages and rising prices of conventional feed such as grains, legumes, etc. for animal production. Moreover, the soaring prices of cereals (e.g., barley, wheat and corn), which are the major energy sources in ruminant diets force nutritionists to search and explore inexpensive alternatives that can partially substitute for the expensive grains. Feeding of unconventional feedstuffs and in some cases agricultural byproducts, which are of no food value to humans, can be one of the solutions to overcome the problem of feed shortages and rising prices.

Cacti (*Opuntia* spp.) has been recognized as one of the most widely used low cost alternative feeds in many parts of the world, especially in semi-arid regions, due to their adaptation to different environmental conditions (Stintzing and Carle 2005; Elghandour et al. 2016c). Cacti has become an important source of green fodder which ensures several livestock species survival in the semi-arid and arid regions of the world with frequent periods of prolonged droughts (Costa et al. 2009). The chemical composition and nutritive value of spineless prickly pear cactus (PC) species differ from region to region depending on many factors including the environment and genotype. The PC is a rich source of non-fibrous carbohydrates (617 g/kg DM), and is an excellent energy source with high dry matter (DM) digestibility (Wanderley et al. 2002). Replacement of energy feedstuff such as corn grain (CG) with PC may require some form of supplementation with other feed additives to improve its fermentation potential and utilization.

Rumen and cecal modifiers such as exogenous fibrolytic enzymes (Kholif et al. 2016a; Morsy et al. 2016), *Saccharomyces cerevisiae* (Rodriguez et al. 2015; Salem et al. 2016a) have been used as ration ingredients for ruminants and horses. Little is known about the nutritive value of *Salix babylonica* (SB) extract in equine nutrition; however, some information is available on ruminant nutrition (Rivero et al. 2016). Extracts of SB have been evaluated as feed additives in ruminant nutrition due to its anti-microbial effects and its ability to modulate ruminal fermentation and improve nutrient utilization (Valdes et al. 2015). The

antimicrobial activity of SB extracts has been attributed to its content of a number of plant secondary metabolites such as alkaloids, saponins and phenolics (Cedillo et al. 2014) which rumen microorganisms have the ability to degrade and utilize as an energy source at low and moderate concentrations without negative effects on rumen fermentation (Salem et al. 2014a, 2016b). In ruminants, SB extract enhanced feed intake (Salem et al. 2014b), daily gain (Cedillo et al. 2014), and milk production (Salem et al. 2014b). It has also been reported to have natural anthelmintic activity (Cedillo et al. 2015; Salem et al. 2016c). To the best of our knowledge, this is the first study to include the extract of SB in the diet of horses. The aim of the current study was to investigate the effects of replacing CG in a horse's diet with PC at different levels in the presence of different levels of *S. babylonica* extract on cecal in vitro gas (GP) and methane (CH₄) production and cecal fermentation kinetics.

Materials and methods

Extract, substrates and treatments

Plant leaves of *S. babylonica* were collected randomly from several young and mature trees during summer of 2015. Leaves were freshly chopped into 1–2 cm lengths and immediately extracted at 1 g leaf/8 mL of water. Plant materials were individually soaked and incubated in water in the laboratory at 25–30 °C for 72 h in jar. After incubation, the jar was heated to 39 °C for 1 h and then immediately filtered and the filtrate collected and stored at 4 °C for further use.

Three total mixed rations were prepared where CG was replaced with PC at three levels (/kg): 0 g (Control), 75 g (PC75) or 150 g (PC150). The extract of SB was added at four levels: 0, 0.6, 1.2 and 1.8 mL/g DM of substrates. The chemical composition and ingredients is shown in Table 1.

In vitro cecal fermentation and biodegradation

Cecal contents (the inoculum source) were collected from 4 Criollo horses (3–4 years of age and weighing 300 ± 15 kg) from the local slaughterhouse of Toluca, Mexico State, Mexico. Horses had about eight hours grazing and were given water twice a day

Table 1 Composition of the experimental diets^a. Adapted from Elghandour et al. 2016b, c

	Control	PC75	PC150
Ingredients (g/kg DM)			
Oats straw	249	248	248
Steam rolled corn	250	175	100
Soybean hulls	250	250	250
Steam rolled barley	120	110	120
Wheat bran	30	30	30
Corn gluten feed	30	30	20
Prickly pear cactus	0	75	150
Molasses	70	80	80
Vitamins/minerals ^b	1	2	2
Chemical composition (g/kg DM)			
Organic matter	964	940	957
Crude protein	130	119	113
Neutral detergent fiber	356	428	340
Acid detergent fiber	121	130	122
Ether extract	24	22	23
Non-structural carbohydrates	455	371	481

^a PC75 prickly pear cactus was included at 75 g/kg DM of total mixed ration; PC150, prickly pear cactus was included at 150 g/kg DM of total mixed ration

^b Contained: Vitamin A (12,000,000 IU), Vitamin D₃ (2,500,000 IU), Vitamin E (15,000 IU), Vitamin K (2.0 g), Vitamin B₁ (2.25 g), Vitamin B₂ (7.5 g), Vitamin B₆ (3.5 g), Vitamin B₁₂ (20 mg), Pantothenic acid (12.5 g), Folic acid (1.5 g), Biotin (125 mg), Niacin (45 g), Fe (50 g), Zn (50 g), Mn (110 g), Cu (12 g), I (0.30 g), Se (200 mg), Co (0.20 g)

without feed supplementation. The horses had grazed predominantly on pasture containing two native grasses (*Festuca arundinacea* and ryegrass). Individual cecal samples were equally collected from the cecum of each horse and then mixed and homogenized to obtain a homogenized sample of fecal contents which were mixed with the Goering and Van Soest (1970) buffer solution without trypticase in the ratio of 1:4 v/v. The incubation media was subsequently mixed and strained through four layers of cheesecloth into a flask with an O₂-free headspace, and used to inoculate three identical runs of incubation in 120-mL serum bottles containing 0.5 g DM of substrate in presence of different doses of SB extract.

Bottles with substrates plus three bottles without substrate and SB as blanks were used. After filling all bottles, they were flushed with CO₂ and immediately closed with rubber stoppers, shaken and placed in an incubator set at 39 °C. Gas production was recorded at

2, 4, 6, 8, 10, 12, 14, 24, 36, 48, 54, 60, and 72 h using the Pressure Transducer Technique (Extech instruments, Waltham, USA) of Theodorou et al. (1994). The production of CH₄ was recorded using Gas-Pro detector (Gas Analyzer CROWCON Model Tetra 3, Abingdon, UK) at 2, 6, 10, 14, 24, 36, 48, 54, 60, and 72 h of incubation.

At the end of incubation after 72 h, bottles were uncapped and the pH was measured using a digital pH meter (Conductronic pH15, Puebla, Mexico), and the residual of each bottle was filtered under vacuum through glass crucibles with a sintered filter and the fermentation residues dried at 65 °C for 72 h to estimate DM disappearance (DMD).

Chemical analyses and calculations

Samples of the rations were analyzed for DM (#934.01), ash (#942.05), N (#954.01) and ether extract (#920.39) according to AOAC (1997) and the ration's contents for neutral detergent fiber content (NDF, Van Soest et al. 1991), acid detergent fiber (ADF) and lignin (AOAC 1997; #973.18) analyses were carried out using an ANKOM²⁰⁰ Fiber Analyzer Unit (ANKOM Technology Corp., Macedon, NY, USA) with the use of an alpha amylase and sodium sulfite.

For estimation of GP kinetic, recorded gas volumes (mL/g DM) were fitted using the NLIN procedure of SAS (2002) according to France et al. (2000) model as:

$$y = b \times [1 - e^{-c(t-Lag)}]$$

where y is the volume of GP at time t (h); b is the asymptotic GP (mL/g DM); c is the fractional rate of fermentation (/h), and Lag (h) is the discrete lag time prior to any gas release.

Metabolizable energy (ME, MJ/kg DM) was estimated according to Menke et al. (1979) as:

$$ME = 2.20 + 0.136 \text{ GP (mL/0.5 g DM)} + 0.057 \text{ CP (g/kg DM)}$$

where GP is net GP in mL from 200 mg of dry sample after 24 h of incubation.

The partitioning factor at 24 h of incubation (PF₂₄; a measure of fermentation efficiency) was calculated as the ratio of DM degradability in vitro (mg) to the volume (mL) of GP at 24 h [i.e., DMD/total GP (GP₂₄)] according to Blümmel et al. (1997). Gas yield (GY₂₄) was calculated as the volume of gas (mL gas/g DM) produced after 24 h of incubation divided by the amount of DMD (g) as:

$GY_{24} = \text{mL gas/g DM/g DMD}$

Short chain fatty acid concentrations (SCFA) were calculated according to Getachew et al. (2002) as:

$SCFA(\text{mmol}/200 \text{ mg DM}) = 0.0222 GP - 0.00425$

where GP is the 24 h net gas production (mL/200 mg DM).

Statistical analyses

Data from each of the three runs within the same sample of each of the three individual samples of rations were averaged prior to statistical analysis, then mean values of each individual sample were used as the experimental unit. Results of in vitro GP and cecal fermentation parameters were analyzed as a factorial experiment using the PROC GLM option of SAS (2002) as:

$Y_{ijk} = \mu + R_i + D_j + (R \times D)_{ij} + E_{ijk}$

where Y_{ijk} = is every observation of the i th ration type (R_i) with j th SB extract dose (D_j); μ is the general mean; $(R \times D)_{ij}$ is the interaction between ration type and SB extract dose; E_{ijk} is the experimental error. Linear and quadratic polynomial contrasts were used to examine responses of different PC rations (levels) to increasing addition levels of SB extract. Statistical significance was declared at $P < 0.05$.

Results

Gas production kinetics

No ration type \times SB extract dose interaction was observed ($P > 0.05$) for all investigated parameters of GP kinetics (Table 2). Ration type affected ($P \leq 0.001$) the asymptotic GP, the rate of GP and the lag time of GP. Ignoring the effect of SB addition (i.e. 0 mL SB/g DM), increasing the level of PC in the ration increased the asymptotic GP (quadratic effect, $P = 0.001$), decreased both of the rate of GP (linear and quadratic effect, $P < 0.001$) and the lag time of GP (quadratic effect, $P < 0.001$). Besides, GP at different incubation hours were affected ($P < 0.05$) by different rations i.e. increasing the level of PC in the ration increased GP values.

The level of SB extract did not affect the asymptotic GP or the rate of GP. Increasing the level of SB extract linearly decreased ($P = 0.001$) the lag time of GP of all diets. However, increasing level of SB extract tended to decreased (quadratic effect, $P = 0.07$) the

rate of gas production. The extract of SB linearly affected ($P < 0.05$) GP at some incubation hours.

Methane production

No interaction was observed ($P > 0.05$) between ration type and SB extract level for CH_4 production. No CH_4 was released before 24 h of incubation (Table 3). With the exception of CH_4 at 36 h of incubation, ration type had no effect ($P > 0.05$) on CH_4 production. The ration with PC75 had the highest ($P = 0.022$) CH_4 production at 36 h of incubation compared with the other rations.

No effect was observed ($P > 0.05$) on CH_4 production with the inclusion of SB extract in the rations at all incubation hours, with the exception of CH_4 at 36 h of incubation; where SB extract addition decreased CH_4 (linear effect, $P = 0.006$; quadratic effect, $P = 0.001$) compared with rations without SB inclusion.

Cecal fermentation kinetics

Interactions were observed between ration type and SB level for cecal pH ($P < 0.001$) and DMD ($P = 0.01$) (Table 4). At the level of 0 mL SB/g DM, the PC75 and PC150 rations increased cecal pH compared with the control ration. The PC150 ration had the highest ($P < 0.05$) DMD, SCFA, and GY_{24} . The inclusion of SB extract had no effect ($P > 0.05$) on all investigated fermentation kinetic parameters.

Discussion

Gas production

No ration type \times SB extract dose interaction was observed ($P > 0.05$) for GP kinetics. Therefore, the effect of ration type and SB extract dose will be individually discussed. Gas production is generally a good indicator of digestibility, fermentability and rumen microbial protein production (Rodriguez et al. 2015; Vallejo et al. 2016). Gas production is dependent on nutrient availability for rumen micro-organisms (Elghandour et al. 2015a, b). The quadratic increases in asymptotic GP and lower rate and lag time of GP with increasing level of PC replacement for CG reveals increased fermentation of the insoluble but

Table 2 In vitro ceal gas kinetics of three levels of prickly pear cactus (PC) at different levels (mg/g DM) of *Salix babylonica* (SB) extract inclusion

Ration ^a	SB extract	Gas production parameters ^b		Gas production (mL/g DM) at:													
		<i>b</i>	<i>c</i>	<i>Lag</i>	2 h	4 h	6 h	8 h	10 h	12 h	14 h	24 h	36 h	48 h	54 h	60 h	72 h
Control	0	381	0.083	8.85	58	107	149	185	214	240	261	329	361	374	377	379	380
	0.6	369	0.125	7.45	80	143	192	231	261	284	303	350	365	368	369	369	369
	1.2	473	0.104	7.32	88	160	218	266	304	335	361	432	461	469	472	471	472
	1.8	463	0.093	7.02	79	144	198	243	280	311	337	412	446	457	459	461	462
PC75	0	414	0.059	8.35	46	86	122	154	183	208	230	310	361	387	394	400	407
	0.6	421	0.072	7.71	54	102	142	177	208	235	258	336	381	402	408	412	417
	1.2	437	0.067	7.34	54	100	141	177	208	235	259	341	391	414	421	425	431
	1.8	472	0.059	3.21	52	98	139	176	208	237	263	355	414	442	451	457	465
PC150	0	548	0.052	3.92	53	102	145	184	219	250	279	384	457	497	509	519	531
	0.6	458	0.059	4.48	51	96	136	172	203	232	257	345	402	430	438	444	451
	1.2	502	0.055	3.34	51	98	139	176	209	239	266	363	428	463	473	481	490
	1.8	538	0.051	2.98	52	99	141	179	213	244	272	376	449	488	501	510	523
Pooled SEM		32.6	0.0067	0.837	5.2	9.3	12.5	15.0	17.0	18.6	19.9	23.9	26.8	28.8	29.6	30.2	31.1
Ration effect																	
	Linear	0.536	<0.001	0.103	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.001	0.014	0.265	0.787	0.991	0.863	0.692
	Quadratic	0.001	<0.001	<0.001	0.001	0.003	0.007	0.017	0.044	0.101	0.210	0.537	0.036	0.005	0.002	0.002	0.001
Dose effect																	
	Linear	0.118	0.643	0.001	0.061	0.057	0.054	0.052	0.049	0.049	0.048	0.050	0.062	0.076	0.083	0.089	0.098
	Quadratic	0.968	0.070	0.644	0.045	0.054	0.065	0.080	0.097	0.117	0.140	0.305	0.538	0.714	0.774	0.820	0.881
	Ration × dose	0.4479	0.448	0.284	0.129	0.132	0.147	0.160	0.171	0.180	0.189	0.196	0.238	0.303	0.363	0.386	0.404

SEM standard error of the mean

^a PC75, prickly pear cactus was included at 75 g/kg DM of total mixed ration; PC150, prickly pear cactus was included at 150 g/kg DM of total mixed ration

^b *b* is the asymptotic gas production (mL/g DM); *c* is the rate of gas production (/h); *Lag* is the initial delay before gas production begins (h)

Table 3 Proportional in vitro methane (CH₄) productions of three levels of prickly pear cactus (PC) at different levels (mg/g DM) of *Salix babylonica* (SB) extract inclusion

Ration ^a	SB extract	Methane production ^b (ml/g DM) at:						
		24 h	36 h	48 h	54 h	60 h	72 h	
Control	0	ND	3.29	3.40	4.65	8.28	8.31	
	0.6	0.84	0.88	0.89	3.08	3.08	4.41	
	1.2	ND	ND	2.03	6.18	6.19	8.04	
	1.8	ND	ND	2.99	4.41	4.99	7.22	
PC75	0	0.82	4.21	4.86	5.65	5.73	6.48	
	0.6	ND	0.53	0.57	1.16	1.17	1.18	
	1.2	ND	0.52	1.15	2.51	2.53	2.56	
	1.8	0.82	4.16	5.83	5.95	7.50	8.17	
PC150	0	1.04	2.76	5.61	5.76	8.54	8.72	
	0.6	1.13	1.29	1.91	4.15	4.19	4.25	
	1.2	0.59	1.16	5.77	5.90	8.82	8.98	
	1.8	ND	0.58	0.64	1.31	1.34	1.38	
Pooled SEM		0.430	0.754	1.346	1.454	2.002	2.162	
Ration effect								
Linear			0.518	0.022	0.422	0.467	0.332	0.130
Quadratic			0.161	0.590	0.362	0.928	0.526	0.980
Dose effect								
Linear			0.332	0.006	0.193	0.230	0.088	0.215
Quadratic			0.422	0.001	0.350	0.815	0.882	0.902
Ration × dose			0.892	0.925	0.050	0.057	0.129	0.144

SEM standard error of the mean, ND not detected (i.e. 0 mL CH₄/g DM)

^a PC75, prickly pear cactus was included at 75 g/kg DM of total mixed ration; PC150, prickly pear cactus was included at 150 g/kg DM of total mixed ration

^b No CH₄ was produced at 2, 6, 10, 14 h of incubation

degradable fraction (Elghandour et al. 2016b). These results suggest a steady increasing availability of carbohydrate fractions to the microbial population growth and activity, in agreement with previous studies (Elghandour et al. 2015a, b). Blümmel and Ørskov (1993) showed that the asymptotic GP can be used to predict feed intake because 88% of variance in intake was due to GP implying the PC75 and PC150 rations had the propensity to induce feed intake and growth rate in cattle (Blümmel and Ørskov 1993). The decreased rate with increasing the level of PC in the ration indicate that the time necessary for degradation was longer than for PC rations, thus less gas was produced in the short term. Microbial growth and accessibility of microbial enzymes to feed particles are reflected by the rate at which different chemical constituents are degraded. Since fractional rate of GP was correlated with feed intake (Khazaal et al. 1995),

PC150 ration would likely enhance feed intake and performance of ruminants. This is because performance is largely a function of feed intake, which is a better indicator of nutritive value of feed than apparent digestibility (Okunade et al. 2014). The discrete lag time prior to GP was decreased with the PC75 and PC150 ration suggesting faster microbial adaptation to the ration, in agreement with previous reports (Elghandour et al. 2015a, b, 2016c).

The inclusion of SB extract did not affect the asymptotic GP or the rate of GP; however, it decreased the lag time of GP. This effect indicates positive effects on ruminal fermentation (Salem et al. 2014a; Elghandour et al. 2015b), and ruminal microorganisms activity possibly due to the ability of rumen microorganisms to degrade secondary metabolites in SB extract and utilize them as an energy source (Hart et al. 2008).

Table 4 Degradation and in vitro cecal fermentation profile of three levels of prickly pear cactus (PC) at different levels (mg/g DM) of *Salix babylonica* (SB) extract inclusion

Ration ^a	SB	pH	ME	DMD	SCFA	PF ₂₄	GY ₂₄
Control	0	5.43	13.1	861	8.3	4.81	208
	0.6	5.10	12.9	930	8.2	4.86	206
	1.2	6.55	15.6	885	10.4	4.66	215
	1.8	6.57	15.3	879	10.1	4.67	214
PC75	0	7.00	13.4	849	8.6	4.79	209
	0.6	6.96	13.8	848	8.9	4.76	210
	1.2	7.01	14.1	861	9.2	4.75	211
	1.8	6.86	14.9	857	9.8	4.70	213
PC150	0	7.07	16.4	877	11.0	4.63	216
	0.6	6.09	14.6	862	9.5	4.72	212
	1.2	6.32	15.4	861	10.2	4.67	214
	1.8	6.19	16.1	859	10.8	4.64	216
Pooled SEM		0.187	0.78	10.5	0.64	0.054	2.3
Ration							
Linear		<0.001	0.776	<0.001	0.784	0.949	0.984
Quadratic		0.894	0.005	0.316	0.005	0.014	0.010
Dose							
Linear		0.796	0.076	0.764	0.076	0.128	0.116
Quadratic		0.415	0.714	0.480	0.714	0.742	0.746
Ration × dose		<0.001	0.363	0.010	0.364	0.486	0.456

DMD, the in vitro dry matter digestibility (mg/g DM); GY₂₄, gas yield at 24 h (mL gas/g DMD); ME is the metabolizable energy (MJ/kg DM); PF₂₄, partitioning factor at 24 h of incubation (mg DMD/mL gas); pH, fermentation pH; SCFA, short-chain fatty acids (mmol/g DM)

SEM standard error of the mean

^a PC75, prickly pear cactus was included at 75 g/kg DM of total mixed ration; PC150, prickly pear cactus was included at 150 g/kg DM of total mixed ration

Methane production

Without the occurrence of ration type × SB extract dose interaction, ration type and SB level did not affect CH₄ production. Methane emission from ruminants depends on diet degradability and chemical composition (Elghandour et al. 2016c, d). Ruminant livestock is one of the sources responsible for greenhouse gas (e.g. CH₄ emission from animal production sector is responsible for about 18% of all greenhouse gas emissions) emission (Intergovernmental Panel on Climate Change 2008). Methane is produced as a result of ruminal fermentation of feed in the rumen causing a loss of digested energy (Johnson and Johnson 1995). With the same rations used in the present experiment, Elghandour et al. (2016c) observed that increasing PC level in the ration linearly

increased the asymptotic CH₄ production; however, in another experiment, the same research team (Elghandour et al. 2016b) observed that replacing CG with soybean hulls did not affect CH₄ production. The different response between soybean hulls and PC may be due to different chemical composition. Besides, the different inoculum source (rumen contents vs cecal contents) can be another reason.

Fermentation kinetics

The rations PC75 and PC150 increased cecal pH. However, Elghandour et al. (2016c) observed a declining pH with increasing level of PC. The difference maybe related with the inoculum source. In their experiment, Elghandour et al. (2016c) used rumen liquor compared with inoculum from cecal

contents in the present experiment. Higher pH is required for the activity of microorganisms presented in the cecum of horses, for their activity and ability to degrade fiber of the diet. The PC150 ration increased DMD, SCFA, and GY_{24} . Feed degradation and fermentation rate has been reported to be directly proportional to GP (Dhanoa et al. 2000). The higher GP is a good indicator of the higher potential degradability of the substrate (i.e. PC150 ration). Higher GP with PC based rations compared to CG ration (control ration) indicates a higher content of highly fermentable constituents of PC than CG. The increased DMD with increasing GP as the level of PC increased confirms the hypothesis that increasing DMD or substrate fermentability ought to be accompanied by increased GP. The improvements of fermentation parameters observed with the replacement of CG by PC could be due to additional availability of the fermentable carbohydrates which possibly promoted microbial growth (Forsberg et al. 2000) and also enhanced the incubation environment. It has been shown that availability of nutrients for rumen microorganisms will stimulate the degradability of different nutrients (Paya et al. 2007). Besides, increased SCFA production and ME are associated with high activities of microbes in the rumen. It can therefore be inferred that PC will supply more fermentable carbohydrates, promote degradability, digestibility and microbial protein synthesis relative to CG. Increasing SCFA with increased PC level was consistent with the increased OMD and ME, in agreement with earlier reports (Elghandour et al. 2013). Increased SCFA is important in terms of enhanced lactose production, milk volume and overall energy balance (Kholif et al. 2015, 2016b). The increases of fermentation profiles with increasing level of PC may be due to increased fiber digestion and enhanced ruminal fermentation (Nsereko et al. 2002) and improved attachment and colonization of PC rations by ruminal micro-organisms (Nsereko et al. 2002; Elghandour et al. 2013). In confirmation of highly positive correlation between ME and GP at 24 h (Menke et al. 1979), both ME and GY_{24} increased with increasing PC level in the ration indicating an improved incubation environment and thus fermentability. The SB extract addition to the ration was not effective in improving all the ruminal fermentation parameters probably due to its

inefficiency in improving fermentation efficiency, fermentation kinetics and GP.

Results reported in the present experiment suggest that prickly pear cactus flour has a potential fermentation efficiency and fermentation profile superior to that of corn grain. It can therefore be used to replace corn grain in concentrate ration to replace conventional energy sources (e.g., maize, barley and sorghum) in ruminant diets. The inclusion of *S. babylonica* extract in the tested rations had a weak effect on their fermentation. The best level of dietary inclusion of prickly pear cactus was 150 g PC/kg DM (replacement of CG at 60%). Further research in which CG is replaced with PC flour with or without SB extract inclusion should be conducted in in vivo trials to validate current findings.

Acknowledgements Authors would like to thank the Universidad Autónoma del Estado de México (project # UAEM 4120/2016SF).

Compliance with ethical standards

Conflict of interest All authors declare that there are no present or potential conflicts of interest among the authors and other people or organizations that could inappropriately bias their work.

References

- Ahmed MH, Elghandour MMY, Salem AZM, Zeweil HS, Kholif AE, Klieve AV, Abdelrassol AMA (2015) Influence of *Trichoderma reesei* or *Saccharomyces cerevisiae* on performance, ruminal fermentation, carcass characteristics and blood biochemistry of lambs fed *Atriplex nummularia* and *Acacia saligna* mixture. *Livest Sci* 180:90–97
- AOAC (1997) Association of official analytical chemists. Official methods of analysis, 16th edn. AOAC, Arlington
- Blümmel M, Ørskov ER (1993) Comparison of in vitro gas production and nylon bag degradability of roughages in predicting feed intake in cattle. *Anim Feed Sci Technol* 40:109–119
- Blümmel M, Steingss H, Becker K (1997) The relationship between in vitro gas production, in vitro microbial biomass yield and ^{15}N incorporation and its implications for the prediction of voluntary feed intake of roughages. *Br J Nutr* 77:911–921
- Cedillo J, Vázquez-Armijo JF, González-Reyna A, Salem AZM, Kholif AE, Hernández-Meléndez J, Martínez-González JC, de Oca Jiménez RM, Rivero N, López D (2014) Effects of different doses of *Salix babylonica* extract on growth performance and diet in vitro gas production in Pelibuey growing lambs. *Ital J Anim Sci* 13(3):609–613

- Cedillo J, Kholif AE, Salem AZM, Elghandour MMY, Vázquez JF, Alonso MU, Barbabosa A, Chagoyán JCV, Reyna AG (2015) Oral administration of Sauce llorón extract to growing lambs to control gastrointestinal nematodes and *Moniezia* spp. *Asian Pac J Trop Med* 8(7):520–525
- Costa RG, Filho EMB, de Medeiros AN, Givisiez PEN, Queirogac RCRE, Silva Melo AAS (2009) Effects of increasing levels of cactus pear (*Opuntia ficus-indica* L. Miller) in the diet of dairy goats and its contribution as a source of water. *Small Rumin Res* 82:62–65
- Dhanoa MS, Lopez S, Dijkstra J (2000) Estimating the extent of degradation of ruminant feeds from a description of their gas production profiles observed in vitro: comparison of models. *Br J Nutr* 83:131–142
- Elghandour MMY, Salem AZM, Gonzalez-Ronquillo M, Bórquez JL, Gado HM, Odongo NE, Penueles CG (2013) Effects of exogenous enzymes on in vitro gas production kinetics and ruminal fermentation of four fibrous feeds. *Anim Feed Sci Technol* 179:46–53
- Elghandour MMY, Kholif AE, Bastida AZ, Martínez DLP, Salem AZM (2015a) *In vitro* gas production of five rations of different maize silage and concentrate ratios influenced by increasing levels of chemically characterized extract of *Salix babylonica*. *Turk J Vet Anim Sci* 39(2):186–194
- Elghandour MMY, Kholif AE, Marquez-Molina O, Vazquez-Armijo JF, Puniya AK, Salem AZM (2015b) Influence of individual or mixed cellulase and xylanase mixture on in vitro rumen gas production kinetics of total mixed rations with different maize silage and concentrate ratios. *Turk J Vet Anim Sci* 39(4):435–442
- Elghandour MMY, Kholif AE, Hernández J, Mariezcurrena MD, López S, Camacho LM, Márquez O, Salem AZM (2016a) Influence of the addition of exogenous xylanase with or without pre-incubation on the in vitro ruminal fermentation of three fibrous feeds. *Czech J Anim Sci* 61(6):262–272
- Elghandour MMY, Kholif AE, Salem AZM, de Oca RM, Barbabosa A, Mariezcurrena M, Olafadehan OA (2016b) Addressing sustainable ruminal methane and carbon dioxide emissions of soybean hulls by organic acid salts. *J Clean Prod* 135:194–200
- Elghandour MMY, Kholif AE, Salem AZM, Olafadehan OA, Kholif AM (2016c) Sustainable anaerobic rumen methane and carbon dioxide productions from prickly pear cactus flour by organic acid salts addition. *J Clean Prod* 139:1362–1369
- Elghandour MMY, Vázquez JC, Salem AZM, Kholif AE, Cipriano MM, Camacho LM, Márquez O (2016d) *In vitro* gas and methane production of two mixed rations influenced by three different cultures of *Saccharomyces cerevisiae*. *J Appl Anim Res* 45(1):389–395
- Forsberg C, Forano E, Chesson A (2000) Microbial adherence to the plant cell wall and enzymatic hydrolysis. In: Cronje PB (ed) *Ruminant physiology: digestion, metabolism, growth and reproduction*. CABI Publishing, Wallingford, pp 79–97
- France J, Dijkstra J, Dhanoa MS, López S, Bannink A (2000) Estimating the extent of degradation of ruminant feeds from a description of their gas production profiles observed in vitro: derivation of models and other mathematical considerations. *Br J Nutr* 83:143–150
- Getachew G, Makkar HPS, Becker K (2002) Tropical browses: contents of phenolic compounds, in vitro gas production and stoichiometric relationship between short chain fatty acid and in vitro gas production. *J Agric Sci Camb* 139:341–352
- Goering MK, Van Soest PJ (1970) Forage fiber analysis (apparatus, reagents, procedures and some applications). *Agriculture handbook*, No. 379. Agricultural Research Service, USDA, Washington, DC
- Hart KJ, Yanez-Ruiz DR, Duval SM, McEwan NR, Newbold CJ (2008) Plant extracts to manipulate rumen fermentation. *Anim Feed Sci Technol* 147:8–35
- Intergovernmental Panel on Climate Change (2008) *Climate change, 2007. Synthesis report. Contribution of working group I, II, and III to the fourth assessment report of the intergovernmental panel on climate change*. IPCC, Geneva
- Johnson KA, Johnson DE (1995) Methane emissions from cattle. *J Anim Sci* 73:2483–2492
- Khazaal K, Dentinho MT, Ribeiro JM, Ørskov ER (1995) Prediction of apparent digestibility and voluntary feed intake of hays fed to sheep: comparison between using fibre component, in vitro digestibility or characteristics of gas production or nylon bag degradation. *Anim Sci* 61:521–538
- Kholif AE, Gouda GA, Morsy TA, Salem AZM, Lopez S, Kholif AM (2015) *Moringa oleifera* leaf meal as a protein source in lactating goat's diets: feed intake, digestibility, ruminal fermentation, milk yield and composition, and its fatty acids profile. *Small Rumin Res* 129:129–137
- Kholif AE, Baza-García LA, Elghandour MMY, Salem AZM, Barbabosa A, Dominguez-Vara IA, Sanchez-Torres JE (2016a) In vitro assessment of fecal inocula from horses fed on high-fiber diets with fibrolytic enzymes addition on gas, methane, and carbon dioxide productions as indicators of hindgut activity. *J Equine Vet Sci* 39:44–50
- Kholif AE, Morsy TA, Abd El Tawab AM, Anele UY, Galyean ML (2016b) Effect of supplementing diets of Anglo-Nubian goats with soybean and flaxseed oils on lactational performance. *J Agric Food Chem* 64(31):6163–6170
- Menke KH, Raab L, Salewski A, Steingass H, Fritz D, Schneider W (1979) The estimation of the digestibility and metabolizable energy content of ruminant feeding stuffs from the gas production when they are incubated with rumen liquor in vitro. *J Agric Sci Camb* 93:217–222
- Morsy TA, Kholif AE, Kholif SM, Kholif AM, Sun X, Salem AZM (2016) Effects of two enzyme feed additives on digestion and milk production in lactating Egyptian buffaloes. *Ann Anim Sci* 16:209–222
- Nsereko VL, Beauchemin KA, Morgavi DP, Rode LM, Furtado AF, McAllister TA, Iwaasa AD, Yang WZ, Wang Y (2002) Effect of a fibrolytic enzyme preparation from *Trichoderma longibrachiatum* on the rumen microbial population of dairy cows. *Can J Microbiol* 48:14–20
- Okunade SA, Olafadehan OA, Isah OA (2014) Fodder potential and acceptability of selected tree leaves by goats. *Anim Nutr Feed Technol* 14:489–498
- Paya H, Taghizadeh A, Janmohammadi H, Moghadam GA (2007) Nutrient digestibility and gas production of some tropical feeds used in ruminant diets estimated by the in vivo and in vitro gas production techniques. *Am J Anim Vet Sci* 2:108–113

- Rivero N, Salem AZM, Ayala M, Elghandour MMY, Kholif AE, Barbabosa A, Camacho LM, Rojas S, Olivares J, Cipriano M (2016) Influence of *Salix babylonica* extract, exogenous enzyme of xylanase and their combination on blood hematological and biochemical profile in sheep and goats. *Indian J Anim Sci* 86(10):1140–1144
- Rodriguez MP, Mariezcurrena MD, Mariezcurrena MA, Lagunas BC, Elghandour MMY, Kholif AM, Kholif AE, Almaraz EM, Salem AZM (2015) Influence of live cells or cells extract of *Saccharomyces cerevisiae* on in vitro gas production of a total mixed ration. *Ital J Anim Sci* 14:590–595
- Salem AZM, Kholif AE, Elghandour MMY, Buendía G, Mariezcurrena MD, Hernandez SR, Camacho LM (2014a) Influence of oral administration of *Salix babylonica* extract on milk production and composition in dairy cows. *Ital J Anim Sci* 13(1):10–14
- Salem AZM, Ryena AC, Elghandour MMY, Camacho LM, Kholif AE, Salazar MC, Domínguez IA, Jiménez RM, Almaraz EM, Martínez AGL, Mariezcurrena MA (2014b) Influence of *Salix babylonica* extract in combination or not with increasing levels of minerals mixture on in vitro rumen gas production kinetics of a total mixed ration. *Ital J Anim Sci* 13:873–879
- Salem AZ, Elghandour MM, Kholif AE, Barbabosa A, Camacho LM, Odongo NE (2016a) Influence of feeding horses a high fiber diet with or without live yeast cultures supplementation on feed intake, nutrient digestion, blood chemistry, fecal coliform count, and in vitro fecal fermentation. *J Equine Vet Sci* 39:12–19
- Salem AZM, Elghandour MMY, Kholif AE, López S, Pliego AB, Cipriano-Salazar M, Chagoyán JCV, de Oca Jiménez RM, Alonso MU (2016b) Tree leaves of *Salix babylonica* extract as a natural anthelmintic for small-ruminant farms in a semiarid region in Mexico. *Agrofor Syst*. doi:10.1007/s10457-016-9909-z
- Salem AZM, Kholif AE, Elghandour MMY, Hernández J, Limas A, Cipriano M, Camacho L, Rojas S, Olivares J (2016c) Influence of *Salix babylonica* extract addition on in vitro rumen gas production and degradability of ryegrass silage harvested in different cutting days. *Indian J Anim Sci* 86(9):1030–1035
- SAS (2002) Statistical analysis system. User's guide: statistics. Ver 9.0. SAS Institute, Cary
- Stintzing G, Carle R (2005) Cactus stems (*Opuntia* spp.): a review on their chemistry and uses. *Mol Nutr Food Res* 49:175
- Theodorou MK, Williams BA, Dhanoa MS, McAllan AB, France J (1994) A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. *Anim Feed Sci Technol* 48:185–197
- Valdes KI, Salem AZM, López S, Alonso MU, Rivero N, Elghandour MMY, Domínguez IA, Ronquillo MG, Kholif AE (2015) Influence of exogenous enzymes in presence of *Salix babylonica* extract on digestibility, microbial protein synthesis and performance of lambs fed maize silage. *J Agric Sci Camb* 153:732–742
- Vallejo L, Salem AZM, Kholif AE, Elghangour M, Fajardo R, Rivero N, Bastida A, Mariezcurrena M (2016) Influence of cellulase or xylanase on the in vitro rumen gas production and fermentation of corn stover. *Indian J Anim Sci* 86(1):70–74
- Van Soest PJ, Robertson JB, Lewis BA (1991) Methods for dietary fibre, neutral detergent fibre, and non-starch carbohydrates in relation to animal nutrition. *J Dairy Sci* 74:3583–3597
- Wanderley WL, Ferreira MA, Andrade DKB, Vêras ASC, Farias I, Lima LE, Dias AMA (2002) Replacement of forage cactus (*Opuntia ficus indica* Mill) for sorghum silage (*Sorghum bicolor* (L.) Moench) in the dairy cows feeding. *Rev Bras Zootec* 31:273–281